

REGULARITY THEORY FOR 2-DIMENSIONAL ALMOST MINIMAL CURRENTS III: BLOWUP

CAMILLO DE LELLIS, EMANUELE SPADARO AND LUCA SPOLAOR

ABSTRACT. We analyze the asymptotic behavior of a 2-dimensional integral current which is almost minimizing in a suitable sense at a singular point. Our analysis is the second half of an argument which shows the discreteness of the singular set for the following three classes of 2-dimensional currents: area minimizing in Riemannian manifolds, semi-calibrated and spherical cross sections of 3-dimensional area minimizing cones.

This paper is the fourth and last in a series of works aimed at establishing an optimal regularity theory for 2-dimensional integral currents which are almost minimizing in a suitable sense. Building upon the monumental work of Almgren [1], Chang in [3] established that 2-dimensional area-minimizing currents in Riemannian manifolds are classical minimal surfaces, namely they are regular (in the interior) except for a discrete set of branching singularities. The argument of Chang is however not entirely complete since a key starting point of his analysis, the existence of the so-called “branched center manifold”, is only sketched in the appendix of [3] and requires the understanding (and a suitable modification) of the most involved portion of the monograph [1].

An alternative proof of Chang’s theorem has been found by Rivière and Tian in [13] for the special case of J -holomorphic curves. Later on the approach of Rivière and Tian has been generalized by Bellettini and Rivière in [2] to handle a case which is not covered by [3], namely that of special Legendrian cycles in S^5 (see also [?] for a further generalization).

Meanwhile the first and second author revisited Almgren’s theory giving a much shorter version of his program for proving that area-minimizing currents are regular up to a set of Hausdorff codimension 2, cf. [4, 5, 6, 7, 8]. In this note and its companion papers [10, 9] we build upon the latter works in order to give a complete regularity theory which includes both the theorems of Chang and Bellettini-Rivière as special cases. In order to be more precise, we introduce the following terminology (cf. [11, Definition 0.3]).

Definition 0.1. Let $\Sigma \subset \mathbb{R}^{m+n}$ be a C^2 submanifold and $U \subset \mathbb{R}^{m+n}$ an open set.

- (a) An m -dimensional integral current T with finite mass and $\text{spt}(T) \subset \Sigma \cap U$ is area-minimizing in $\Sigma \cap U$ if $\mathbf{M}(T + \partial S) \geq \mathbf{M}(T)$ for any $m + 1$ -dimensional integral current S with $\text{spt}(S) \subset \subset \Sigma \cap U$.
- (b) A semicalibration (in Σ) is a C^1 m -form ω on Σ such that $\|\omega_x\|_c \leq 1$ at every $x \in \Sigma$, where $\|\cdot\|_c$ denotes the comass norm on $\Lambda^m T_x \Sigma$. An m -dimensional integral current T with $\text{spt}(T) \subset \Sigma$ is *semicalibrated* by ω if $\omega_x(\vec{T}) = 1$ for $\|T\|$ -a.e. x .
- (c) An m -dimensional integral current T supported in $\partial \mathbf{B}_{\bar{R}}(p) \subset \mathbb{R}^{m+n}$ is a *spherical cross-section of an area-minimizing cone* if $p \ast T$ is area-minimizing.

In what follows, given an integer rectifiable current T , we denote by $\text{Reg}(T)$ the subset of $\text{spt}(T) \setminus \text{spt}(\partial T)$ consisting of those points x for which there is a neighborhood U such that $T \llcorner U$ is a (constant multiple of) a regular submanifold. Correspondingly, $\text{Sing}(T)$ is the set $\text{spt}(T) \setminus (\text{spt}(\partial T) \cup \text{Reg}(T))$. Observe that $\text{Reg}(T)$ is relatively open in $\text{spt}(T) \setminus \text{spt}(\partial T)$ and thus $\text{Sing}(T)$ is relatively closed. The main result of this and the works [10, 9] is then the following

Theorem 0.2. *Let $m = 2$ and T be as in (a), (b) or (c) of Definition 0.1. Assume in addition that Σ is of class C^{3,ε_0} (in case (a) and (b)) and ω of class C^{2,ε_0} (in case (b)) for some positive ε_0 . Then $\text{Sing}(T)$ is discrete.*

Clearly Chang's result is covered by case (a). As for the case of special Lagrangian cycles considered by Bellettini and Rivière in [2] observe that they form a special subclass of both (b) and (c). Indeed these cycles arise as spherical cross-sections of 3-dimensional special lagrangian cones: as such they are then spherical cross sections of area-minimizing cones but they are also semicalibrated by a specific smooth form on \mathbb{S}^5 .

Following the Almgren-Chang program, Theorem 0.2 will be established through a suitable "blow-up argument": this argument is presented here but requires several tools. The first important tool is the theory of multiple-valued functions, for which we will use the results and terminology of the papers [4, 5]. The second tool is a suitable approximation result for area-minimizing currents with graphs of multiple valued functions, which for the case at hand has been established in the preceding note [10]. The last tool is the so-called "branched center manifold": this has been constructed in the paper [9]. We note in passing that all our arguments use heavily the uniqueness of tangent cones for T . This result is a, by now classical, theorem of White for area-minimizing 2-dimensional currents in the euclidean space, cf. [15]. Chang extended it to case (a) in the appendix of [3], whereas Pumberger and Rivière covered case (b) in [12]. A general derivation of these results for a wide class of almost minimizers has been given in [11]: the theorems in there cover, in particular, all the cases of Definition 0.1.

The proof of Theorem 0.2 is based, as in [3], on an induction statement, cf. Theorem 1.8 below. Although Theorem 1.8 is already stated in [9], we will recall it in the next section, where we also show how it implies Theorem 0.2. This and the previous paper [9] can be hence thought as the two halves in the proof of Theorem 1.8. After introducing some terminology, in Section 2 we will state the first half in Theorem 2.6 (which is proved in [9]) and the second half in Theorem 2.8, and we will then show how they fit together to prove Theorem 1.8. The remaining sections (the biggest portion of this note) are then dedicated to prove Theorem 2.8.

Acknowledgments The research of Camillo De Lellis and Luca Spolaor has been supported by the ERC grant RAM (Regularity for Area Minimizing currents), ERC 306247.

1. PRELIMINARIES AND THE MAIN INDUCTION STATEMENT

1.1. Basic notation and first main assumptions. For the notation concerning submanifolds $\Sigma \subset \mathbb{R}^{2+n}$ we refer to [6, Section 1]. With $\mathbf{B}_r(p)$ and $B_r(x)$ we denote, respectively, the open ball with radius r and center p in \mathbb{R}^{2+n} and the open ball with radius r

and center x in \mathbb{R}^2 . $\mathbf{C}_r(p)$ and $\mathbf{C}_r(x)$ will always denote the cylinder $B_r(x) \times \mathbb{R}^n$, where $p = (x, y) \in \mathbb{R}^2 \times \mathbb{R}^n$. We will often need to consider cylinders whose bases are parallel to other 2-dimensional planes, as well as balls in m -dimensional affine planes. We then introduce the notation $B_r(p, \pi)$ for $\mathbf{B}_r(p) \cap (p + \pi)$ and $\mathbf{C}_r(p, \pi)$ for $B_r(p, \pi) + \pi^\perp$. e_i will denote the unit vectors in the standard basis, π_0 the (oriented) plane $\mathbb{R}^2 \times \{0\}$ and $\vec{\pi}_0$ the 2-vector $e_1 \wedge e_2$ orienting it. Given an m -dimensional plane π , we denote by \mathbf{p}_π and \mathbf{p}_π^\perp the orthogonal projections onto, respectively, π and its orthogonal complement π^\perp . For what concerns integral currents we use the definitions and the notation of [14]. Since π is used recurrently for 2-dimensional planes, the 2-dimensional area of the unit circle in \mathbb{R}^2 will be denoted by ω_2 .

By [10, Lemma 1.1] in case (b) we can assume, without loss of generality, that the ambient manifold Σ coincides with the euclidean space \mathbb{R}^{2+n} . In the rest of the paper we will therefore always make the following

Assumption 1.1. T is an integral current of dimension 2 with bounded support and it satisfies one of the three conditions (a), (b) or (c) in Definition 0.1. Moreover

- In case (a), $\Sigma \subset \mathbb{R}^{2+n}$ is a C^{3,ε_0} submanifold of dimension $2 + \bar{n} = 2 + n - l$, which is the graph of an entire function $\Psi : \mathbb{R}^{2+\bar{n}} \rightarrow \mathbb{R}^l$ and satisfies the bounds

$$\|D\Psi\|_0 \leq c_0 \quad \text{and} \quad \mathbf{A} := \|A_\Sigma\|_0 \leq c_0, \quad (1.1)$$

where c_0 is a positive (small) dimensional constant and $\varepsilon_0 \in]0, 1[$.

- In case (b) we assume that $\Sigma = \mathbb{R}^{2+n}$ and that the semicalibrating form ω is C^{2,ε_0} .
- In case (c) we assume that T is supported in $\Sigma = \partial\mathbf{B}_R(p_0)$ for some p_0 with $|p_0| = R$, so that $0 \in \partial\mathbf{B}_R(p_0)$. We assume also that $T_0\partial\mathbf{B}_R(p_0)$ is \mathbb{R}^{2+n-1} (namely $p_0 = (0, \dots, 0, \pm|p_0|)$) and we let $\Psi : \mathbb{R}^{2+n-1} \rightarrow \mathbb{R}$ be a smooth extension to the whole space of the function which describes Σ in $\mathbf{B}_2(0)$. We assume then that (1.1) holds, which is equivalent to the requirement that R^{-1} be sufficiently small.

In addition, since the conclusion of Theorem 0.2 is local, by [11, Proposition 0.4] we can also assume to be always in the following situation.

Assumption 1.2. In addition to Assumption 1.1 we assume the following:

- $\partial T \llcorner \mathbf{C}_2(0, \pi_0) = 0$;
- $0 \in \text{spt}(T)$ and the tangent cone at 0 is given by $\Theta(T, 0) \llbracket \pi_0 \rrbracket$ where $\Theta(T, 0) \in \mathbb{N} \setminus \{0\}$;
- T is irreducible in any neighborhood U of 0 in the following sense: it is not possible to find S, Z non-zero integer rectifiable currents in U with $\partial S = \partial Z = 0$ (in U), $T = S + Z$ and $\text{spt}(S) \cap \text{spt}(Z) = \{0\}$.

In order to justify point (iii), observe that we can argue as in the proof of [11, Theorem 3.1]: assuming that in a certain neighborhood U there is a decomposition $T = S + Z$ as above, it follows from [11, Proposition 2.2] that both S and Z fall in one of the classes of Definition 0.1. In turn this implies that $\Theta(S, 0), \Theta(Z, 0) \in \mathbb{N} \setminus \{0\}$ and thus $\Theta(S, 0) < \Theta(T, 0)$. We can then replace T with either S or Z . Let $T_1 = S$ and argue similarly if it is not irreducible: obviously we can apply one more time the argument above and find a

T_2 which satisfies all the requirements and has $0 < \Theta(T_2, 0) < \Theta(T_1, 0)$. This process must stop after at most $N = \Theta(T, 0)$ steps: the final current is then necessarily irreducible.

1.2. Branching model. We next introduce an object which will play a key role in the rest of our work, because it is the basic local model of the singular behavior of a 2-dimensional area-minimizing current: for each positive natural number Q we will denote by $\mathfrak{B}_{Q,\rho}$ the flat Riemann surface which is a disk with a conical singularity, in the origin, of angle $2\pi Q$ and radius $\rho > 0$. More precisely we have

Definition 1.3. $\mathfrak{B}_{Q,\rho}$ is topologically an open 2-dimensional disk, which we identify with the topological space $\{(z, w) \in \mathbb{C}^2 : w^Q = z, |z| < \rho\}$. For each $(z_0, w_0) \neq 0$ in $\mathfrak{B}_{Q,\rho}$ we consider the connected component $\mathfrak{D}(z_0, w_0)$ of $\mathfrak{B}_{Q,\rho} \cap \{(z, w) : |z - z_0| < |z_0|/2\}$ which contains (z_0, w_0) . We then consider the smooth manifold given by the atlas

$$\{(\mathfrak{D}(z, w), (x_1, x_2)) : (z, w) \in \mathfrak{B}_{Q,\rho} \setminus \{0\}\},$$

where (x_1, x_2) is the function which gives the real and imaginary part of the first complex coordinate of a generic point of $\mathfrak{B}_{Q,\rho}$. On such smooth manifold we consider the following flat Riemannian metric: on each $\mathfrak{D}(z, w)$ with the chart (x_1, x_2) the metric tensor is the usual euclidean one $dx_1^2 + dx_2^2$. Such metric will be called the *canonical flat metric* and denoted by e_Q . The coordinates $(x_1, x_2) = z$ will be called *standard flat coordinates*.

When $Q = 1$ we can extend smoothly the metric tensor to the origin and we obtain the usual euclidean 2-dimensional disk. For $Q > 1$ the metric tensor does not extend smoothly to 0, but we can nonetheless complete the induced geodesic distance on $\mathfrak{B}_{Q,\rho}$ in a neighborhood of 0: for $(z, w) \neq 0$ the distance to the origin will then correspond to $|z|$. The resulting metric space is a well-known object in the literature, namely a flat Riemann surface with an isolated conical singularity at the origin (see for instance [16]). Note that for each z_0 and $0 < r \leq \min\{|z_0|, \rho - |z_0|\}$ the set $\mathfrak{B}_{Q,\rho} \cap \{|z - z_0| < r\}$ consists then of Q nonintersecting 2-dimensional disks, each of which is a geodesic ball of $\mathfrak{B}_{Q,\rho}$ with radius r and center (z_0, w_i) for some $w_i \in \mathbb{C}$ with $w_i^Q = z_0$. We then denote each of them by $B_r(z_0, w_i)$ and treat it as a standard disk in the euclidean 2-dimensional plane (which is correct from the metric point of view). We use however the same notation for the distance disk $B_r(0)$, namely for the set $\{(z, w) : |z| < r\}$, although the latter is *not isometric* to the standard euclidean disk. Since this might be create some ambiguity, we will use the specification $\mathbb{R}^2 \supset B_r(0)$ when referring to the standard disk in \mathbb{R}^2 .

1.3. Admissible Q -branchings. When one of (or both) the parameters Q and ρ are clear from the context, the corresponding subscript (or both) will be omitted. We will consider repeatedly functions u defined on \mathfrak{B} . We will always treat each point of \mathfrak{B} as an element of \mathbb{C}^2 , mostly using z and w for the horizontal and vertical complex coordinates. Often \mathbb{C} will be identified with \mathbb{R}^2 and thus the coordinate z will be treated as a two-dimensional real vector, avoiding the more cumbersome notation (x_1, x_2) .

Definition 1.4 (Q -branchings). Let $\alpha \in]0, 1[$, $b > 1$, $Q \in \mathbb{N} \setminus \{0\}$ and $n \in \mathbb{N} \setminus \{0\}$. An admissible α -smooth and b -separated Q -branching in \mathbb{R}^{2+n} (shortly a Q -branching) is the

graph

$$\text{Gr}(u) := \{(z, u(z, w)) : (z, w) \in \mathfrak{B}_{Q, 2\rho}\} \subset \mathbb{R}^{2+n} \quad (1.2)$$

of a map $u : \mathfrak{B}_{Q, 2\rho} \rightarrow \mathbb{R}^n$ satisfying the following assumptions. For some constants $C_i > 0$ we have

- u is continuous, $u \in C^{3, \alpha}$ on $\mathfrak{B}_{Q, \rho} \setminus \{0\}$ and $u(0) = 0$;
- $|D^j u(z, w)| \leq C_i |z|^{1-j+\alpha} \forall (z, w) \neq 0$ and $j \in \{0, 1, 2, 3\}$;
- $[D^3 u]_{\alpha, B_r(z, w)} \leq C_i |z|^{-2}$ for every $(z, w) \neq 0$ with $|z| = 2r$;
- If $Q > 1$, then there is a positive constant $c_s \in]0, 1[$ such that

$$\min\{|u(z, w) - u(z, w')| : w \neq w'\} \geq 4c_s |z|^b \quad \text{for all } (z, w) \neq 0. \quad (1.3)$$

The map $\Phi(z, w) := (z, u(z, w))$ will be called the *graphical parametrization* of the Q -branching.

Any Q -branching as in the Definition above is an immersed disk in \mathbb{R}^{2+n} and can be given a natural structure as integer rectifiable current, which will be denoted by \mathbf{G}_u . For $Q = 1$ a map u as in Definition 1.4 is a (single valued) $C^{1, \alpha}$ map $u : \mathbb{R}^2 \supset B_{2\rho}(0) \rightarrow \mathbb{R}^n$. Although the term branching is not appropriate in this case, the advantage of our setup is that $Q = 1$ will not be a special case in the induction statement of Theorem 1.8 below. Observe that for $Q > 1$ the map u can be thought as a Q -valued map $u : \mathbb{R}^2 \supset B_\rho(0) \rightarrow \mathcal{A}_Q(\mathbb{R}^n)$, setting $u(z) = \sum_{(z, w_i) \in \mathfrak{B}} \llbracket u(z, w_i) \rrbracket$ for $z \neq 0$ and $u(0) = Q \llbracket 0 \rrbracket$. The notation $\text{Gr}(u)$ and \mathbf{G}_u is then coherent with the corresponding objects defined in [5] for general Q -valued maps.

1.4. The inductive statement. Before coming to the key inductive statement, we need to introduce some more terminology.

Definition 1.5 (Horned Neighborhood). Let $\text{Gr}(u)$ be a b -separated Q -branching. For every $a > b$ we define the *horned neighborhood* $\mathbf{V}_{u, a}$ of $\text{Gr}(u)$ to be

$$\mathbf{V}_{u, a} := \{(x, y) \in \mathbb{R}^2 \times \mathbb{R}^n : \exists (x, w) \in \mathfrak{B}_{Q, 2\rho} \text{ with } |y - u(x, w)| < c_s |x|^a\}, \quad (1.4)$$

where c_s is the constant in (1.3).

Definition 1.6 (Excess). Given an m -dimensional current T in \mathbb{R}^{m+n} with finite mass, its *excess* in the ball $\mathbf{B}_r(x)$ and in the cylinder $\mathbf{C}_r(p, \pi')$ with respect to the m -plane π are

$$\mathbf{E}(T, \mathbf{B}_r(p), \pi) := (2\omega_m r^m)^{-1} \int_{\mathbf{B}_r(p)} |\vec{T} - \vec{\pi}|^2 d\|T\| \quad (1.5)$$

$$\mathbf{E}(T, \mathbf{C}_r(p, \pi'), \pi) := (2\omega_m r^m)^{-1} \int_{\mathbf{C}_r(p, \pi')} |\vec{T} - \vec{\pi}|^2 d\|T\|. \quad (1.6)$$

For cylinders we omit the third entry when $\pi = \pi'$, i.e. $\mathbf{E}(T, \mathbf{C}_r(p, \pi)) := \mathbf{E}(T, \mathbf{C}_r(p, \pi), \pi)$. In order to define the spherical excess we consider T as in Assumption 1.1 and we say that π *optimizes the excess* of T in a ball $\mathbf{B}_r(x)$ if

- In case (b)

$$\mathbf{E}(T, \mathbf{B}_r(x)) := \min_{\tau} \mathbf{E}(T, \mathbf{B}_r(x), \tau) = \mathbf{E}(T, \mathbf{B}_r(x), \pi); \quad (1.7)$$

- In case (a) and (c) $\pi \subset T_x \Sigma$ and

$$\mathbf{E}(T, \mathbf{B}_r(x)) := \min_{\tau \subset T_x \Sigma} \mathbf{E}(T, \mathbf{B}_r(x), \tau) = \mathbf{E}(T, \mathbf{B}_r(x), \pi). \quad (1.8)$$

Note in particular that, in case (a) and (c), $\mathbf{E}(T, \mathbf{B}_r(x))$ differs from the quantity defined in [8, Definition 1.1], where, although Σ does not coincide with the ambient euclidean space, τ is allowed to vary among *all* planes, as in case (b). Thus a notation more consistent with that of [8] would be, in case (a) and (c), $\mathbf{E}^\Sigma(T, \mathbf{B}_r(x))$. However, the difference is a minor one and we prefer to keep our notation simpler.

Our main induction assumption is then the following

Assumption 1.7 (Inductive Assumption). T is as in Assumption 1.1 and 1.2. For some constants $\bar{Q} \in \mathbb{N} \setminus \{0\}$ and $0 < \alpha < \frac{1}{2\bar{Q}}$ there is an α -admissible \bar{Q} -branching $\text{Gr}(u)$ with $u : \mathfrak{B}_{\bar{Q},2} \rightarrow \mathbb{R}^n$ such that

- (Sep) If $\bar{Q} > 1$, u is b -separated for some $b > 1$; a choice of some $b > 1$ is fixed also in the case $\bar{Q} = 1$, although in this case the separation condition is empty.
- (Hor) $\text{spt}(T) \subset \mathbf{V}_{u,a} \cup \{0\}$ for some $a > b$;
- (Dec) There exist $\gamma > 0$ and a $C_i > 0$ with the following property. Let $p = (x_0, y_0) \in \text{spt}(T) \cap \mathbf{C}_{\sqrt{2}}(0)$ and $4d := |x_0| > 0$, let V be the connected component of $\mathbf{V}_{u,a} \cap \{(x, y) : |x - x_0| < d\}$ containing p and let $\pi(p)$ be the plane tangent to $\text{Gr}(u)$ at the only point of the form $(x_0, u(x_0, w_i))$ which is contained in V . Then

$$\mathbf{E}(T \llcorner V, \mathbf{B}_\sigma(p), \pi(p)) \leq C_i^2 d^{2\gamma-2} \sigma^2 \quad \forall \sigma \in [\frac{1}{2}d^{(b+1)/2}, d]. \quad (1.9)$$

The main inductive step is then the following theorem, where we denote by $T_{p,r}$ the rescaled current $(\iota_{p,r})_\# T$, through the map $\iota_{p,r}(q) := (q - p)/r$.

Theorem 1.8 (Inductive statement). *Let T be as in Assumption 1.7 for some $\bar{Q} = Q_0$. Then,*

- (a) *either T is, in a neighborhood of 0, a Q multiple of a \bar{Q} -branching $\text{Gr}(v)$;*
- (b) *or there are $r > 0$ and $Q_1 > Q$ such that $T_{0,r}$ satisfies Assumption 1.7 with $\bar{Q} = Q_1$.*

Theorem 0.2 follows then easily from Theorem 1.8 and [11].

1.5. Proof of Theorem 0.2. As already mentioned, without loss of generality we can assume that Assumption 1.1 holds, cf. [11, Lemma 1.1] (the bounds on \mathbf{A} and Ψ can be achieved by a simple scaling argument). Fix now a point p in $\text{spt}(T) \setminus \text{spt}(\partial T)$. Our aim is to show that T is regular in a punctured neighborhood of p . Without loss of generality we can assume that p is the origin. Upon suitably decomposing T in some neighborhood of 0 we can easily assume that (I) in Assumption 1.7 holds, cf. the argument of Step 4 in the proof of [11, Theorem 3.1]. Thus, upon suitably rescaling and rotating T we can assume that π_0 is the unique tangent cone to T at 0, cf. [11, Theorem 3.1]. In fact, by [11, Theorem 3.1] T satisfies Assumption 1.7 with $\bar{Q} = 1$: it suffices to chose $u \equiv 0$ as admissible smooth branching. If T were not regular in any punctured neighborhood of 0, we could then apply Theorem 1.8 inductively to find a sequence of rescalings T_{0,ρ_j} with $\rho_j \downarrow 0$ which satisfy Assumption 1.7 with $\bar{Q} = Q_j$ for some strictly increasing sequence of

integers. It is however elementary that the density $\Theta(0, T)$ bounds Q_j from above, which is a contradiction.

2. THE BRANCHED CENTER MANIFOLD AND THE BLOW-UP THEOREM

From now on we fix T satisfying Assumption 1.7. Observe that, without loss of generality, we are always free to rescale homothetically our current T with a factor larger than 1 and ignore whatever portion falls outside $\mathbf{C}_2(0)$. We will do this several times, with factors which will be assumed to be sufficiently large. Hence, if we can prove that something holds in a sufficiently small neighborhood of 0, then we can assume, without loss of generality, that it holds on \mathbf{C}_2 . For this reason we can assume that the constants C_i in Definition 1.4 and Assumption 1.7 is as small as we want. In turns this implies that there is a well-defined orthogonal projection $\mathbf{P} : \mathbf{V}_{u,a} \cap \mathbf{C}_1 \rightarrow \text{Gr}(u) \cap \mathbf{C}_2$, which is a $C^{2,\alpha}$ map. We next recall [9, Lemma 2.1]:

Lemma 2.1. *Let T and u be as in Assumption 1.7 for some \bar{Q} . Then the nearest point projection $\mathbf{P} : \mathbf{V}_{u,a} \cap \mathbf{C}_1 \rightarrow \text{Gr}(u)$ is a well-defined $C^{2,\alpha}$ map. In addition there is $Q \in \mathbb{N} \setminus \{0\}$ such that $\Theta(0, T) = Q\bar{Q}$ and the unique tangent cone to T at 0 is $Q\bar{Q} \llbracket \pi_0 \rrbracket$. Finally, after possibly rescaling T , $\Theta(p, T) \leq Q$ for every $p \in \mathbf{C}_2 \setminus \{0\}$ and, for every $x \in B_2(0)$, each connected component of $(x \times \mathbb{R}^n) \cap \mathbf{V}_{u,a}$ contains at least one point of $\text{spt}(T)$.*

Since we will assume during the rest of the paper that the above discussion applies, we summarize the relevant conclusions in the following

Assumption 2.2. *T satisfies Assumption 1.7 for some \bar{Q} and with C_i sufficiently small. $Q \geq 1$ is an integer, $\Theta(0, T) = Q\bar{Q}$ and $\Theta(p, T) \leq Q$ for all $p \in \mathbf{C}_2 \setminus \{0\}$.*

The overall plan to prove Theorem 1.8 is then the following:

- (CM) We construct first a branched center manifold, i.e. a second admissible smooth branching φ on $\mathfrak{B}_{\bar{Q}}$, and a corresponding Q -valued map N defined on the normal bundle of $\text{Gr}(\varphi)$, which approximates T with a very high degree of accuracy (in particular more accurately than u) and whose average $\eta \circ N$ is very small;
- (BU) Assuming that alternative (a) in Theorem 1.8 does not hold, we study the asymptotic behavior of N around 0 and use it to build a new admissible smooth branching v on some $\mathfrak{B}_{k\bar{Q}}$ where $k \geq 2$ is a factor of Q : this map will then be the one sought in alternative (b) of Theorem 1.8 and a suitable rescaling of T will lie in a horned neighborhood of its graph.

The first part of the program is the one achieved in [9], whereas the second part will be completed in this note. Note that, when $Q = 1$, from (BU) we will conclude that alternative (a) necessarily holds: this will be a simple corollary of the general case, but we observe that it could also be proved resorting to the classical Allard's regularity theorem.

2.1. Smallness condition. In several occasions we will need that the ambient manifold Σ is suitably flat and that the excess of the current T is suitably small. This can, however, be easily achieved after scaling. More precisely we recall [9, Lemma 2.3].

Lemma 2.3. *Let T be as in the Assumptions 1.7 and 2.2. After possibly rescaling, rotating and modifying Σ outside $\mathbf{C}_2(0)$ we can assume that, in case (a) and (c) of Definition 0.1,*

- (i) Σ is a complete submanifold of \mathbb{R}^{2+n} ;
- (ii) $T_0\Sigma = \mathbb{R}^{2+\bar{n}} \times \{0\}$ and, $\forall p \in \Sigma$, Σ is the graph of a C^{3,ε_0} map $\Psi_p : T_p\Sigma \rightarrow (T_p\Sigma)^\perp$.

Under these assumptions, we denote by \mathbf{c} and \mathbf{m}_0 the following quantities

$$\mathbf{c} := \sup\{\|D\Psi_p\|_{C^{2,\varepsilon_0}} : p \in \Sigma\} \quad \text{in the cases (a) and (c) of Definition 0.1} \quad (2.1)$$

$$\mathbf{c} := \|d\omega\|_{C^{1,\varepsilon_0}} \quad \text{in case (b) of Definition 0.1} \quad (2.2)$$

$$\mathbf{m}_0 := \max\{\mathbf{c}^2, \mathbf{E}(T, \mathbf{C}_2, \pi_0), C_i^2, c_s^2\}, \quad (2.3)$$

where C_i and c_s are the constants appearing in Definition 1.4 and Assumption 1.7. Then, for any $\varepsilon_2 > 0$, after possibly rescaling the current by a large factor, we can assume

$$\mathbf{m}_0 \leq \varepsilon_2. \quad (2.4)$$

In order to carry on the plan outlined in the previous subsection, it is convenient to use a different parametrization of Q -branchings. If we remove the origin, any admissible Q -branching is a Riemannian submanifold of $\mathbb{R}^{2+n} \setminus \{0\}$: this gives a Riemannian tensor $g := \Phi^\sharp e$ (where e denotes the euclidean metric on \mathbb{R}^{2+n}) on the punctured disk $\mathfrak{B}_{Q,2\rho} \setminus \{0\}$. Note that in (z, w) the difference between the metric tensor g and the canonical flat metric e_Q can be estimated by (a constant times) $|z|^{2\alpha}$: thus, as it happens for the canonical flat metric e_Q , when $Q > 1$ it is not possible to extend the metric g to the origin. However, using well-known arguments in differential geometry, we can find a conformal map from $\mathfrak{B}_{Q,r}$ onto a neighborhood of 0 which maps the conical singularity of $\mathfrak{B}_{Q,r}$ in the conical singularity of the Q -branching. In fact, we need the following accurate estimates for such a map, cf. [9, Proposition 2.4]:

Proposition 2.4 (Conformal parametrization). *Given an admissible b -separated α -smooth Q -branching $\text{Gr}(u)$ with $\alpha < 1/(2Q)$ there exist a constant $C_0(Q, \alpha) > 0$, a radius $r > 0$ and functions $\Psi : \mathfrak{B}_{Q,r} \rightarrow \text{Gr}(u)$ and $\lambda : \mathfrak{B}_{Q,r} \rightarrow \mathbb{R}_+$ such that*

- (i) Ψ is a homeomorphism of $\mathfrak{B}_{Q,r}$ with a neighborhood of 0 in $\text{Gr}(u)$;
- (ii) $\Psi \in C^{3,\alpha}(\mathfrak{B}_{Q,r} \setminus \{0\})$, with the estimates

$$|D^l(\Psi(z, w) - (z, 0))| \leq C_0 C_i |z|^{1+\alpha-l} \quad \text{for } l = 0, \dots, 3, z \neq 0, \quad (2.5)$$

$$[D^3\Psi]_{\alpha, B_r(z,w)} \leq C_0 C_i |z|^{-2} \quad \text{for } z \neq 0 \text{ and } r = |z|/2; \quad (2.6)$$

- (iii) Ψ is a conformal map with conformal factor λ , namely, if we denote by e the ambient euclidean metric in \mathbb{R}^{2+n} and by e_Q the canonical euclidean metric of $\mathfrak{B}_{Q,r}$,

$$g := \Psi^\sharp e = \lambda e_Q \quad \text{on } \mathfrak{B}_{Q,r} \setminus \{0\}. \quad (2.7)$$

- (iv) The conformal factor λ satisfies

$$|D^l(\lambda - 1)(z, w)| \leq C_0 C_i |z|^{2\alpha-l} \quad \text{for } l = 0, 1, \dots, 2 \quad (2.8)$$

$$[D^2\lambda]_{\alpha, B_r(z,w)} \leq C_0 C_i |z|^{\alpha-2} \quad \text{for } z \neq 0 \text{ and } r = |z|/2. \quad (2.9)$$

Definition 2.5. A map Ψ as in Proposition 2.4 will be called a *conformal parametrization* of an admissible Q -branching.

2.2. The center manifold and the approximation. We are now ready to state the two “halves” of Theorem 1.8. The first one is [9, Theorem 2.6], which we recall here for the reader’s convenience.

Theorem 2.6 (Center Manifold Approximation). *Let T be as in Assumptions 1.7 and 2.2. Then there exist $\eta_0, \gamma_0, r_0, C > 0, b > 1$, an admissible b -separated γ_0 -smooth \bar{Q} -branching \mathcal{M} , a corresponding conformal parametrization $\Psi : \mathfrak{B}_{\bar{Q},2} \rightarrow \mathcal{M}$ and a Q -valued map $\mathcal{X} : \mathfrak{B}_{\bar{Q},2} \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+n})$ with the following properties:*

(i) $\bar{Q}Q = \Theta(T, 0)$ and

$$|D(\Psi(z, w) - (z, 0))| \leq C \mathbf{m}_0^{1/2} |z|^{\gamma_0} \quad (2.10)$$

$$|D^2 \Psi(z, w)| + |z|^{-1} |D^3 \Psi(z, w)| \leq C \mathbf{m}_0^{1/2} |z|^{\gamma_0-1}; \quad (2.11)$$

in particular, if we denote by $A_{\mathcal{M}}$ the second fundamental form of $\mathcal{M} \setminus \{0\}$,

$$|A_{\mathcal{M}}(\Psi(z, w))| + |z|^{-1} |D_{\mathcal{M}} A_{\mathcal{M}}(\Psi(z, w))| \leq C \mathbf{m}_0^{1/2} |z|^{\gamma_0-1}.$$

(ii) $\mathcal{X}_i(z, w)$ is orthogonal to the tangent plane, at $\Psi(z, w)$, to \mathcal{M} .

(iii) If we define $S := T_{0,r_0}$, then $\text{spt}(S) \cap \mathbf{C}_1 \setminus \{0\}$ is contained in a suitable horned neighborhood of the \bar{Q} -branching, where the orthogonal projection \mathbf{P} onto it is well-defined. Moreover, for every $r \in]0, 1[$ we have

$$\|\mathcal{X}|_{B_r}\|_0 + \sup_{p \in \text{spt}(S) \cap \mathbf{P}^{-1}(\Psi(B_r))} |p - \mathbf{P}(p)| \leq C \mathbf{m}_0^{1/4} r^{1+\gamma_0/2}. \quad (2.12)$$

(iv) If we define

$$\mathbf{D}(r) := \int_{B_r} |D\mathcal{X}|^2 \quad \text{and} \quad \mathbf{H}(r) := \int_{\partial B_r} |\mathcal{X}|^2,$$

$$\mathbf{F}(r) := \int_0^r \frac{\mathbf{H}(t)}{t^{2-\gamma_0}} dt \quad \text{and} \quad \mathbf{\Lambda}(r) := \mathbf{D}(r) + \mathbf{F}(r),$$

then the following estimates hold for every $r \in]0, 1[$:

$$\text{Lip}(\mathcal{X}|_{B_r}) \leq C \min\{\mathbf{\Lambda}^{\eta_0}(r), \mathbf{m}_0^{\eta_0} r^{\eta_0}\} \quad (2.13)$$

$$\mathbf{m}_0^{\eta_0} \int_{B_r} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{X}(z, w)| \leq C \mathbf{\Lambda}^{\eta_0}(r) \mathbf{D}(r) + C \mathbf{F}(r). \quad (2.14)$$

(v) Finally, if we set

$$\mathcal{F}(z, w) := \sum_i \llbracket \Psi(z, w) + \mathcal{X}_i(z, w) \rrbracket,$$

then

$$\|S - \mathbf{T}_{\mathcal{F}}(\mathbf{P}^{-1}(\Psi(B_r)))\| \leq C \mathbf{\Lambda}^{\eta_0}(r) \mathbf{D}(r) + C \mathbf{F}(r). \quad (2.15)$$

The second main step is the analysis of the asymptotic behaviour of \mathcal{N} around the origin, which is the main focus of this paper.

Remark 2.7. In order to state it, we agree to define $W^{1,2}$ functions on \mathfrak{B} in the following fashion: removing the origin 0 from \mathfrak{B} we have a C_{loc}^3 (flat) Riemannian manifold embedded in \mathbb{R}^4 and we can define $W^{1,2}$ maps on it following [4]. Alternatively we can use the conformal parametrization $\mathbf{W} : \mathbb{R}^2 = \mathbb{C} \rightarrow \mathfrak{B}_{\bar{Q}}$ given by $\mathbf{W}(z) = (z^{\bar{Q}}, z)$ and agree that $u \in W^{1,2}(\mathfrak{B}_{\bar{Q}})$ if $u \circ \mathbf{W}$ is in $W^{1,2}(\mathbb{R}^2)$. Since discrete sets have zero 2-capacity, it is immediate to verify that these two definitions are equivalent.

In a similar fashion, we will ignore the origin when integrating by parts Lipschitz vector fields, treating $\mathfrak{B}_{\bar{Q}}$ as a C^1 Riemannian manifold. It is straightforward to show that our assumption is correct, for instance removing a disk of radius ε centered at the origin, integrating by parts and then letting $\varepsilon \downarrow 0$.

Theorem 2.8 (Blowup Analysis). *Under the assumptions of Theorem 2.6, the following dichotomy holds:*

- (i) *either there exists $s > 0$ such that $\mathcal{N}|_{B_s} \equiv Q \llbracket 0 \rrbracket$;*
- (ii) *or there exist constants $I_0 > 1$, $a_0, \bar{r}, C > 0$ and an I_0 -homogeneous nontrivial Dir-minimizing function $g : \mathfrak{B}_{\bar{Q}} \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+n})$ such that*
 - $\boldsymbol{\eta} \circ g \equiv 0$,
 - $g = \sum_i \llbracket (0, \bar{g}_i, 0) \rrbracket$, where $\bar{g}_i(x) \in \mathbb{R}^{\bar{n}}$ and $(0, \bar{g}_i(x), 0) \in \mathbb{R}^2 \times \mathbb{R}^{\bar{n}} \times \mathbb{R}^l$,
 - *and the following estimates hold:*

$$\mathcal{G}(\mathcal{N}(z, w), g(z, w)) \leq C|z|^{I_0+a_0} \quad \forall (z, w) \in \mathfrak{B}_Q, |z| < \bar{r}, \quad (2.16)$$

$$\int_{B_{r+\rho} \setminus B_{r-\rho}} |D\mathcal{N}|^2 \leq C r^{2I_0+a_0} + C r^{2I_0-1} \rho \quad \forall 4\rho \leq r < 1, \quad (2.17)$$

$$\mathbf{H}(r) \leq C r \mathbf{D}(r) \quad \forall r < 1. \quad (2.18)$$

Remark 2.9. Note that, when $\bar{Q} = \Theta(T, 0)$, we necessarily have $Q = 1$ and the second alternative is excluded. In particular we conclude that T coincides with $\llbracket \mathcal{M} \rrbracket$ in a neighborhood of 0 and thus it is a regular submanifold in a *punctured neighborhood* of 0.

Remark 2.10. By a simple dyadic argument it follows from (2.17) and (2.18) that

$$\int_{B_r} |D\mathcal{N}|^2 \leq C r^{2I_0} \quad \text{and} \quad \mathbf{F}(r) \leq C r^{2I_0+\gamma_0} \quad \forall r < 1. \quad (2.19)$$

Below we show how to conclude Theorem 1.8 from Theorem 2.8. The remaining part of the paper is dedicated to the proof of the latter, which will be split in six sections each corresponding to one of the following steps.

- (i) In Section 3 we will deduce an almost minimizing property for the map \mathcal{N} in terms of its Dirichlet energy.
- (ii) In Section 4 we will apply the almost minimizing property and compare the Dirichlet energy of \mathcal{N} with that of a suitable harmonic extension of its boundary value on any given ball.

- (iii) In Section 5 we use the comparison above and a first variation argument to derive a suitable Poincaré-type inequality for \mathcal{N} .
- (iv) In Section 6 we compute again the first variations of the Dirichlet energy of \mathcal{N} and use the Poincaré inequality to bound efficiently several error terms.
- (v) Using the latter bounds, in Section 7 we will prove an almost monotonicity property for the frequency function and the existence and boundedness of its limit, which is indeed the number I_0 of Theorem 2.8. The almost minimality of \mathcal{N} will then allow us to conclude an exponential rate of decay to this limit.
- (vi) From the decay of the previous step we will capture in Section 8 the asymptotic behaviour of \mathcal{N} and show the existence of the map g of Theorem 2.8.

The overall strategy follows the ideas and some of the computations in [3]. However several adjustments are needed to carry on the proof in the cases (b) and (c) of Definition 0.1. In particular in Section 7 we need to introduce a suitable modification of the usual frequency function to handle case (b).

2.3. Proof of the inductive step. In the next sections we will prove Theorem 2.8. We start observing that if case (a) of Theorem 1.8 does not hold, then we are necessarily in case (ii) of Theorem 2.8. Therefore we only need to prove that Theorem 2.8(ii) implies Theorem 1.8(b).

We divide the proof in different steps.

Step 1. For a reason which will become clear later, it is convenient to slightly modify the map g to a multivalued map $n(z, w) = \sum_i \llbracket n_i(z, w) \rrbracket$ in such a way that $n_i(z, w)$ is orthogonal to \mathcal{M} at $\Psi(z, w)$. To achieve this it suffices to project $g_i(z, w) = (0, \bar{g}_i(z, w), 0)$ on the normal bundle. Observe that, by the estimates on $|A_{\mathcal{M}}|$ and Ψ , we have

$$|g_i(z, w) - n_i(z, w)| \leq CC_i |z|^{\gamma_0} |g_i(z, w)|, \quad (2.20)$$

$$|Dn|(z, w) \leq |Dg|(z, w) + CC_i |z|^{\gamma_0-1} |g|(z, w). \quad (2.21)$$

We introduce the function $H : \mathfrak{B}_{\bar{Q}} \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+n})$ given by

$$H(z, w) = \sum_{i=1}^Q \llbracket H_i(z, w) \rrbracket := \sum_{i=1}^Q \llbracket \Psi(z, w) + n_i(z, w) \rrbracket .$$

Note that, since g is I_0 -homogeneous and Dir-minimizer, by [4, Proposition 5.1] there is a constant C such that

$$|g_i(z, w) - g_j(z, w)| \geq 2C |z|^{I_0} \quad \text{whenever } g_i(z, w) \neq g_j(z, w). \quad (2.22)$$

In fact [4, Proposition 5.1] is stated for maps with domain $\mathbf{C} = \mathbb{R}^2$. However, if we define the map $\mathbf{W} : \mathbb{C} \rightarrow \mathfrak{B}_{\bar{Q}, \infty}$ as $\mathbf{W}(z) = (z^{\bar{Q}}, z)$, by the conformality of \mathbf{W} it is easy to check that $g \circ \mathbf{W}$ is Dir-minimizer and $I_0 Q$ homogeneous.

By (2.20) and (2.22), provided z is small enough we have

$$|H_i(z, w) - H_j(z, w)| \geq C |z|^{I_0} \quad \text{whenever } H_i(z, w) \neq H_j(z, w). \quad (2.23)$$

Let $\bar{a} \in]0, a_0[$ be a constant to be fixed momentarily and $\zeta := I_0 + \bar{a}/2 > 1$. Set

$$\mathbf{V}_{H,\zeta} := \{H_i(z, w) + p \in \mathbb{R}^{2+n} : |p| < |z|^\zeta, i = 1, \dots, Q\}.$$

We claim that there exists $s > 0$ such that $\text{spt}(T) \cap \mathbf{B}_s \setminus \{0\} \subset \mathbf{V}_{H,\zeta}$.

In order to prove this claim, we distinguish two cases. First we consider any point $p \in \text{spt}(T) \cap \text{spt}(\mathbf{T}_{\mathcal{F}}) \setminus \{0\}$. In this case $p = \Psi(z, w) + \mathcal{N}_i(z, w)$ for some $(z, w) \in \mathfrak{B}_{\bar{Q}} \setminus \{0\}$ and for some $i = 1, \dots, Q$. Without loss of generality, by (2.16) we can assume $|\mathcal{N}_i(z, w) - g_i(z, w)| \leq C|z|^{I_0 + \bar{a}}$, i.e.

$$\begin{aligned} |p - H_i(z, w)| &= |\mathcal{N}_i(z, w) - n_i(z, w)| \leq |\mathcal{N}_i(z, w) - g_i(z, w)| + |g_i(z, w) - n_i(z, w)| \\ &\leq C|z|^{I_0 + \bar{a}} + C|z|^{\gamma_0 + I_0}, \end{aligned} \quad (2.24)$$

which in particular implies $\text{spt}(T) \cap \text{spt}(\mathbf{T}_{\mathcal{F}}) \cap \mathbf{B}_s \subset \mathbf{V}_{H,\zeta}$ if s is sufficiently small and we impose $\frac{\bar{a}}{2} < \gamma_0$.

For the second case we consider a point $p \in \text{spt}(T) \setminus \text{spt}(\mathbf{T}_{\mathcal{F}})$ and assume by contradiction that $p \notin \mathbf{V}_{H,\zeta}$. In particular, in view of (2.24) we have that

$$B := \mathbf{B}_{\frac{|z|^\zeta}{2}}(p) \cap \text{spt}(\mathbf{T}_{\mathcal{F}}) = \emptyset \quad (2.25)$$

if $|z|$ is sufficiently small. By the monotonicity formula we know that $\|T\|(B) \geq C|z|^{2\zeta}$; nevertheless since $B \subset \mathbf{P}^{-1}(B_{2|z|} \setminus B_{|z|/2})$, (2.25) implies $\|T\|(B) \leq \|T - \mathbf{T}_{\mathcal{F}}\|(B)$ and from (2.15) and (2.19) we conclude $\|T\|(B) \leq C|z|^{2I_0 + 2\kappa}$ with $\kappa = \min\{2\eta_0 I_0, \gamma_0\}$. This gives a contradiction if $\bar{a} < 2\kappa$.

Step 2. From the previous step we can infer that g is a constant multiple of an irreducible function, namely there exists $Q' > 0$ such that $\text{card}(g(z, w)) = Q'$ for every $(z, w) \neq (0, 0)$ and there exists a continuous map $h : \mathfrak{B}_{\bar{Q}Q'} \rightarrow \mathbb{R}^{2+n}$ such that

$$g(z, w) = \frac{Q}{Q'} \sum_{\tilde{z}=z, \tilde{w}^{Q'}=w} [[h(\tilde{z}, \tilde{w})]]. \quad (2.26)$$

If this is not the case, by [4, Proposition 5.1] we can decompose g in the superposition of irreducible functions, i.e. there exists a unique decomposition $g = \sum_{j=1}^J k_j g_j$ where $g_j : \mathfrak{B}_Q \rightarrow \mathcal{A}_{q_j}(\mathbb{R}^n)$ are Dir-minimizing I_0 -homogeneous functions, for some choice of positive integers J, k_j, q_j such that $\sum_{j=1}^J k_j q_j = Q$.

Denoting by H^j the corresponding maps

$$H^j(z, w) := \sum_{l=1}^{q_j} [[\Psi(z, w) + (n^j)_l(z, w)]]$$

and by $\mathbf{V}_{H^j,\zeta}$ the corresponding horned neighborhoods

$$\mathbf{V}_{H^j,\zeta} := \{(H^j)_l(z, w) + p \in \mathbb{R}^{2+n} : |p| < |z|^\zeta, l = 1, \dots, q_j\},$$

it follows from (2.23) that the closures of the V_{ζ, H^i} intersect only at the origin. Setting $T_i := T \llcorner V_{\zeta, H^i}$, we infer that $T = \sum_i T_i$ with $\text{spt}(T_i) \cap \text{spt}(T_j) = \{0\}$, against the irreducibility of T . Note that, since $\boldsymbol{\eta} \circ g = 0$ it also follows that $Q' > 1$.

Having established (2.26), let us define $\Theta : \mathfrak{B}_{\bar{Q}Q'} \rightarrow \mathbb{R}^{2+n}$ as

$$\Theta(\tilde{z}, \tilde{w}) := \Psi(\tilde{z}, \tilde{w}^{Q'}) + h^n(\tilde{z}, \tilde{w}) \quad \forall (\tilde{z}, \tilde{w}) \in \mathfrak{B}_{\bar{Q}Q'},$$

where $h^n(\tilde{z}, \tilde{w})$ is the projection of $h(\tilde{z}, \tilde{w})$ on the space normal to \mathcal{M} at the point $\Psi(\tilde{z}, \tilde{w}^{Q'})$. It follows that $\text{Im}(H) = \text{Im}(\Theta)$ is an admissible $\bar{Q}Q'$ -branching (the Hölder regularity for the graphical parametrization follows from the fact that $I_0 > 1$). Moreover, from the homogeneity of g we easily infer that $\text{Im}(\Theta)$ is I_0 -separated (for a suitable constant c_s). Note that for $\zeta' := I_0 + \bar{a}/4$ and s sufficiently small $\mathbf{V}_{H,\zeta} \cap \mathbf{B}_s \subset \mathbf{V}_{\Theta,\zeta'} \cap \mathbf{B}_s$.

Step 3. Finally we prove the condition (Dec) of Assumption 1.7. Let $(z, w) \in \mathfrak{B}_{\bar{Q}}$ with $0 < |z| < \sqrt{2}$, let V be the connected component of $\mathbf{V}_{\Theta,\zeta'} \cap \{(x, y) : |x - z| < d\}$ with $d := |z|/4$ containing $\Theta(z, w)$, and $p \in \text{spt}(T) \cap V$ with coordinates $p = (z, y)$. Denote by π the oriented two-vector for $\text{Im}(\Theta)$ at $\Theta(z, w)$, and consider $\rho \in [\frac{1}{2}d^{(I_0+1)/2}, d]$.

Since $\mathbf{B}_\rho(p) \cap \text{spt}(T) \subset \mathbf{P}^{-1}(\Psi(B_{|z|+2\rho} \setminus B_{|z|-2\rho}))$, we start estimating as follows

$$\begin{aligned} \int_{\mathbf{B}_\rho(p)} |\vec{T} - \vec{\pi}|^2 d\|T \llcorner V\| &\leq \int_{\mathbf{B}_\rho(p) \cap V} |\vec{\mathbf{T}}_{\mathcal{F}} - \vec{\pi}|^2 d\|\mathbf{T}_{\mathcal{F}}\| + \|T - \mathbf{T}_{\mathcal{F}}\|(\mathbf{P}^{-1}(B_{|x_0|+2\rho})) \\ &\stackrel{(2.15)}{\leq} \int_{\mathbf{B}_\rho(p) \cap V} |\vec{\mathbf{T}}_{\mathcal{F}} - \vec{\pi}|^2 d\|\mathbf{T}_{\mathcal{F}}\| + C|z|^{2I_0+2\kappa}. \end{aligned} \quad (2.27)$$

Next, note that for $|z|$ small enough $\mathbf{P}(\mathbf{B}_\rho(p) \cap \mathbf{V}_{\Theta,\zeta'}) \subset \Psi(B_{2\rho}(z, w))$.

We can consider the set of indices $A \subset \{1, \dots, Q\}$ such that $\mathcal{F}_i(z, w) \in V$ for $i \in A$ and estimate as follows

$$\begin{aligned} \int_{\mathbf{B}_\rho(p) \cap V} |\vec{\mathbf{T}}_{\mathcal{F}} - \vec{\pi}|^2 d\|\mathbf{T}_{\mathcal{F}}\| &\leq C \sum_{i \in A} \int_{B_{2\rho}(z, w)} |\vec{\mathbf{T}}_{\mathcal{F}_i(\zeta, \omega)} - \vec{\mathbf{T}}_{\Theta(\zeta, \omega)}|^2 d\zeta + C\rho^2 \text{Lip}(D\Theta|_{B_{2\rho}(z, w)})^2 \\ &\leq C \sum_{i \in A} \int_{B_{2\rho}(z, w)} |\vec{\mathbf{T}}_{\mathcal{F}_i(\zeta, \omega)} - \vec{\mathbf{T}}_{\Psi(\zeta, \omega)}|^2 d\zeta \\ &\quad + C \int_{B_{2\rho}(z, w)} |\vec{\mathbf{T}}_{\Psi(\zeta, \omega)} - \vec{\mathbf{T}}_{\Theta(\zeta, \omega)}|^2 d\zeta + C\rho^4 |z|^{2\theta-2}, \end{aligned} \quad (2.28)$$

where $\theta := \min\{\gamma_0, I_0 - 1\}$ and we used that $|D^2\Theta|(z, w) \leq C|z|^{\theta-1}$.

We can finally use the computation of the excess in curvilinear coordinates in [5, Proposition 3.4] to get

$$\begin{aligned} \sum_i \int_{B_{2\rho}(z, w)} |\vec{\mathbf{T}}_{\mathcal{F}_i(\zeta, \omega)} - \vec{\mathbf{T}}_{\Psi(\zeta, \omega)}|^2 &\leq C \int_{B_{2\rho}(z, w)} (|D\mathcal{N}|^2 + |\zeta|^{2\gamma_0-2} |\mathcal{N}|^2) \\ &\stackrel{(2.19)}{\leq} C \int_{B_{|z|+2\rho} \setminus B_{|z|-2\rho}} |D\mathcal{N}|^2 + C|z|^{2I_0+2\gamma_0} \end{aligned} \quad (2.29)$$

$$\stackrel{(2.17)}{\leq} C|z|^{2I_0+a_0} + C|z|^{2I_0-1} \rho, \quad (2.30)$$

and similarly

$$\begin{aligned}
\int_{B_{2\rho}(z,w)} |\vec{\mathbf{T}}_{\Theta(\zeta,\omega)} - \vec{\mathbf{T}}_{\Psi(\zeta,\omega)}|^2 &\leq C \int_{B_{2\rho}(z,w)} (|Dn|^2 + |\zeta|^{2\gamma_0-2}|n|^2) \\
&\leq C \int_{B_{2\rho}(z,w)} (|Dg|^2 + |\zeta|^{2\gamma_0-2}|g|^2) \\
&\leq C |z|^{2I_0-2}\rho^2 + C |z|^{2I_0+2\gamma_0}
\end{aligned} \tag{2.31}$$

(observe that, in order to apply [5, Proposition 3.4] we need that n takes value into the normal bundle).

Collecting all the estimates together, we have that there exists a suitable constant ϖ such that

$$\int_{\mathbf{B}_\rho(p)} |\vec{T} - \vec{\pi}|^2 d\|T \llcorner V\| \leq C |z|^{2I_0+2\varpi} + C \rho |z|^{2I_0-1} + C \rho^4 |z|^{2\varpi-2} \leq |z|^{\gamma-2} \rho^4, \tag{2.32}$$

where the last inequality is easily verified for $\gamma > 0$ and $|z|$ small enough. This shows (Dec) in Assumption 1.7 and completes the proof.

3. DIRICHLET ALMOST MINIMIZING PROPERTY

The normal approximation \mathcal{N} inherits from T an almost minimizing property for the Dirichlet energy, where the errors involved are in fact expressed in terms of some specific norms of \mathcal{N} itself and of its competitors.

For technical reasons we introduce the maps $F := \sum_{i=1}^Q \llbracket p + N_i(p) \rrbracket$, where $N := \mathcal{N} \circ \Psi^{-1}$. In order to state the almost minimizing property of \mathcal{N} we introduce an appropriate notion of competitor.

Definition 3.1. A Lipschitz map $\mathcal{L}: B_r \rightarrow \mathcal{A}_Q(\mathbb{R}^{n+2})$ is called a competitor for \mathcal{N} in the ball B_r if

- (a) $\mathcal{L}|_{\partial B_r} = \mathcal{N}|_{\partial B_r}$;
- (b) $\text{spt}(\mathcal{G}(z, w)) \subset \Sigma$ for all $(z, w) \in B_r$, where $\mathcal{G}(z, w) := \sum_{j=1}^Q \llbracket \Psi(z, w) + \mathcal{L}_j(z, w) \rrbracket$.

We are now ready to state the almost minimizing property for \mathcal{N} . We use the notation $\mathbf{p}_{T_p\Sigma}$ for the orthogonal projection on the tangent space to Σ at p . We recall that, given our choice of coordinates, $\mathbf{p}_{T_0\Sigma}$ is the projection on $\mathbb{R}^{2+\bar{n}} \times \{0\}$. Since this projection will be used several times, we will denote it by \mathbf{p}_0 . By the $C^{3,\varepsilon}$ regularity of Σ , there exists a map $\Psi_0 \in C^{3,\varepsilon}(\mathbb{R}^{2+\bar{n}}, \mathbb{R}^l)$ such that

$$\Psi_0(0) = 0, \quad D\Psi_0(0) = 0 \quad \text{and} \quad \text{Gr}(\Psi_0) = \Sigma.$$

Next, for each function \mathcal{L} satisfying Condition (b) in Definition 3.1 we consider the map $\bar{\mathcal{L}} := \mathbf{p}_0 \circ \mathcal{L}$, which is a multivalued $\bar{\mathcal{L}}: \mathfrak{B} \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+\bar{n}})$. We observe that it is possible to determine \mathcal{L} from $\bar{\mathcal{L}}$. In particular, fix coordinates $(\xi, \eta) \in \mathbb{R}^{2+\bar{n}} \times \mathbb{R}^l$ and let $\mathcal{L} = \sum \llbracket \mathcal{L}_i \rrbracket$, $\bar{\mathcal{L}} = \sum \llbracket \bar{\mathcal{L}}_i \rrbracket$, where $\bar{\mathcal{L}}_i = \mathbf{p}_0 \circ \mathcal{L}_i$. Then the formula relating \mathcal{L}_i and $\bar{\mathcal{L}}_i$ is

$$\mathcal{L}_i(z, w) = (\bar{\mathcal{L}}_i(z, w), \Psi_0(\mathbf{p}_0(\Psi(z, w)) + \bar{\mathcal{L}}_i(z, w)) - \Psi_0(\mathbf{p}_0(\Psi(z, w)))) . \tag{3.1}$$

Proposition 3.2. *There exists a constant $C_{3.2} > 0$ such that the following holds. If $r \in (0, 1)$ and $\mathcal{L}: B_r \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+n})$ is a Lipschitz competitor for \mathcal{X} with $\|\mathcal{L}\|_\infty \leq r$ and $\text{Lip}(\mathcal{L}) \leq C_{3.2}^{-1}$, then*

$$\int_{B_r} |D\mathcal{X}|^2 \leq (1 + C_{3.2}r) \int_{B_r} |D\bar{\mathcal{L}}|^2 + C_{3.2} \text{Err}_1(\mathcal{X}, B_r) + C_{3.2} \text{Err}_2(\mathcal{L}, B_r) + C_{3.2} r^2 \mathbf{D}'(r), \quad (3.2)$$

where $\bar{\mathcal{L}} := \mathbf{p}_0 \circ \mathcal{L}$ and the error terms $\text{Err}_1(\mathcal{X}, B_r)$, $\text{Err}_2(\mathcal{L}, B_r)$ are given by the following expressions:

$$\text{Err}_1(\mathcal{X}, B_r) = \Lambda^{\eta_0}(r) \mathbf{D}(r) + \mathbf{F}(r) + \mathbf{H}(r) + \mathbf{m}_0^{1/2} r^{1+\gamma_0} \int_{\partial B_r} |\boldsymbol{\eta} \circ \mathcal{X}| \quad (3.3)$$

and

$$\text{Err}_2(\mathcal{L}, B_r) = \mathbf{m}_0^{1/2} \int_{B_r} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{L}|. \quad (3.4)$$

For the proof of Proposition 3.2 we consider separately the three cases:

- (a) T is mass minimizing;
- (b) T is semicalibrated;
- (c) T is the cross-section of a mass minimizing three-dimensional cone.

For notational convenience we set $L := \mathcal{L} \circ \Psi^{-1}$, $G := \mathcal{G} \circ \Psi^{-1}$.

Observe also that, by Lemma A.1 and A.2, it is enough to prove that

$$\int_{B_r} |D\mathcal{X}|^2 \leq (1 + C_{3.2}r) \int_{B_r} |D\mathcal{L}|^2 + C \text{Err}_1(\mathcal{X}, B_r) + C \text{Err}_2(\mathcal{L}, B_r) + \frac{C}{r} \int_{B_r} |\mathcal{L}|^2 + Cr^2 \mathbf{D}'(r). \quad (3.5)$$

Indeed Lemma A.2 implies that

$$\begin{aligned} \int_{B_r} |D\mathcal{L}|^2 &\leq (1 + Cr) \int_{B_r} |D\bar{\mathcal{L}}|^2 + Cr \int_{\partial B_r} |\bar{\mathcal{L}}|^2 \leq (1 + Cr) \int_{B_r} |D\bar{\mathcal{L}}|^2 + Cr \int_{\partial B_r} |\mathcal{L}|^2 \\ &= (1 + Cr) \int_{B_r} |D\bar{\mathcal{L}}|^2 + Cr \int_{\partial B_r} |\mathcal{X}|^2 \leq (1 + Cr) \int_{B_r} |D\bar{\mathcal{L}}|^2 + C \text{Err}_1(\mathcal{X}, B_r), \end{aligned}$$

whereas Lemma A.1 implies

$$\frac{1}{r} \int_{B_r} |\mathcal{L}|^2 \leq Cr \int_{B_r} |D\mathcal{L}|^2 + C \int_{\partial B_r} |\mathcal{L}|^2 \leq Cr \int_{B_r} |D\bar{\mathcal{L}}|^2 + C \text{Err}_1(\mathcal{X}, B_r).$$

3.1. Proof of Proposition 3.2 case (a): T mass minimizing. We fix \mathcal{L} , $\bar{\mathcal{L}}$, L , \mathcal{G} , $\bar{\mathcal{G}}$ and G as above. Let us set

$$Z := T - \mathbf{T}_{\mathcal{J}|_{B_r}} + \mathbf{T}_{\mathcal{G}}. \quad (3.6)$$

Since $\mathcal{F}|_{\partial B_r} = \mathcal{G}|_{\partial B_r}$, from [5] it follows that $\partial(\mathbf{T}_{\mathcal{G}} - \mathbf{T}_{\mathcal{F}|_{B_r}}) = 0$. Moreover $\text{spt}(Z) \subset \Sigma$ and therefore we must have $\mathbf{M}(T) \leq \mathbf{M}(Z)$. Taking into account (2.15), we conclude that

$$\begin{aligned} \mathbf{M}(\mathbf{T}_{\mathcal{F}|_{B_r}}) &\leq \mathbf{M}(T \llcorner \mathbf{p}^{-1}(\Psi(B_r))) + \|T - \mathbf{T}_{\mathcal{F}|_{B_r}}\|(\mathbf{p}^{-1}(\Psi(B_r))) \\ &\leq \mathbf{M}(\mathbf{T}_{\mathcal{G}}) + 2\|T - \mathbf{T}_{\mathcal{F}|_{B_r}}\|(\mathbf{p}^{-1}(\Psi(B_r))) \\ &\leq \mathbf{M}(\mathbf{T}_{\mathcal{G}}) + C \text{Err}_1(\mathcal{X}, B_r). \end{aligned} \quad (3.7)$$

Observe now that $\mathbf{T}_{\mathcal{F}|_{B_r}} = \mathbf{T}_{F|_{\Psi(B_r)}}$ and we can use the Taylor expansion in [5] to compute:

$$\begin{aligned} \mathbf{M}(\mathbf{T}_{\mathcal{F}|_{B_r}}) &\geq Q\mathcal{H}^2(\Psi(B_r)) + \frac{1}{2} \int_{\Psi(B_r)} |DN|^2 - Q \int_{\Psi(B_r)} \langle \boldsymbol{\eta} \circ N, H_{\mathcal{M}} \rangle \\ &\quad - C \int_{\Psi(B_r)} \left(|A_{\mathcal{M}}|^2 |N|^2 + |DN|^4 \right), \end{aligned} \quad (3.8)$$

where $H_{\mathcal{M}}$ denotes the mean curvature vector of \mathcal{M} . Note that in order to apply the Taylor expansion in [5] we need the manifold \mathcal{M} to be C^2 , with an a priori bound on the C^2 norm. However, if we take $T_F \llcorner \mathbf{B}_r \setminus \mathbf{B}_{r/2}$ and rescale by a factor $1/r$, the corresponding rescaled current, map and manifold fall under the assumptions of the Taylor expansion in [5]. We can then scale back to find the corresponding inequalities for $T \llcorner \mathbf{B}_r \setminus \mathbf{B}_{r/2}$ and sum over dyadic annuli to conclude (3.8).

Using the conformality of Ψ we conclude

$$\int_{\Psi(B_r)} |DN|^2 = \int_{B_r} |D\mathcal{X}|^2,$$

As for the other terms, we recall

$$\int_{\Psi(B_r)} |\langle \boldsymbol{\eta} \circ N, H_{\mathcal{M}} \rangle| \leq C\mathbf{m}_0^{1/2} \int_{B_r} |\boldsymbol{\eta} \circ \mathcal{X}| \stackrel{(2.14)}{\leq} C \text{Err}_1(\mathcal{X}, B_r), \quad (3.9)$$

$$\int_{\Psi(B_r)} |DN|^4 \leq C \text{Lip}(\mathcal{X}|_{B_r})^2 \int_{B_r} |D\mathcal{X}|^2 \stackrel{(2.13)}{\leq} C \text{Err}_1(\mathcal{X}, B_r), \quad (3.10)$$

$$\int_{\Psi(B_r)} |A_{\mathcal{M}}|^2 |N|^2 \leq C\mathbf{m}_0 \int_{B_r} |z|^{2\gamma_0-2} |\mathcal{X}|^2 = C\mathbf{m}_0 \int_0^r \frac{\mathbf{H}(s)}{s^{2-2\gamma_0}} ds \leq C \text{Err}_1(\mathcal{X}, B_r). \quad (3.11)$$

Combining the latter estimates with (3.6) and (3.7) we achieve

$$\frac{1}{2} \int_{B_r} |D\mathcal{X}|^2 \leq C \text{Err}_1(\mathcal{X}, B_r) + \mathbf{M}(\mathbf{T}_{\mathcal{G}}) - Q\mathcal{H}^2(\Psi(B_r(x))). \quad (3.12)$$

Next, fix an orthonormal frame ξ_1, ξ_2 on B_r and, using the area formula from [5], compute

$$\begin{aligned} \mathbf{M}(\mathbf{T}_G) &= \int_{\Psi(B_r)} \sum_i |(\xi_1 + DL_i \cdot \xi_1) \wedge (\xi_2 + DL_i \cdot \xi_2)| \\ &\leq \frac{1}{2} \int_{\Psi(B_r)} \sum_i (|\xi_1 + DL_i \cdot \xi_1|^2 + |\xi_2 + DL_i \cdot \xi_2|^2) \\ &= Q\mathcal{H}^2(\Psi(B_r)) + \frac{1}{2} \int_{\Psi(B_r)} |DL|^2 \\ &\quad + Q \int_{\Psi(B_r)} (\langle D(\boldsymbol{\eta} \circ L) \cdot \xi_1, \xi_1 \rangle + \langle D(\boldsymbol{\eta} \circ L) \cdot \xi_2, \xi_2 \rangle). \end{aligned}$$

By conformality the second summand in the last inequality equals $\frac{1}{2} \int_{B_r} |D\mathcal{L}|^2$. We integrate by parts the third summand. Recall that $\boldsymbol{\eta} \circ L = \boldsymbol{\eta} \circ N$ on $\Psi(\partial B_r) = \partial(\Psi(B_r))$: since $\boldsymbol{\eta} \circ N$ is orthogonal to ξ_i the boundary term vanishes. Moreover, since the origin is a singularity, we must in fact integrate by parts in $B_r \setminus B_\varepsilon$ and then let $\varepsilon \rightarrow 0$. A specific choice of ξ_i is $\xi_i = \lambda^{-1/2} D\Psi \cdot e_i$, where e_1, e_2 is the parallel frame on \mathfrak{B}_Q naturally induced by the standard flat coordinates. It then turns out that

$$|D_{\xi_1} \xi_1 + D_{\xi_2} \xi_2|(\Psi(z, w)) \leq C\mathbf{m}_0^{1/2} |z|^{\gamma_0-1}.$$

In particular $|D_{\xi_1} \xi_1 + D_{\xi_2} \xi_2|$ is integrable on B_r and we can therefore conclude

$$\begin{aligned} \mathbf{M}(\mathbf{T}_G) - Q\mathcal{H}^2(\Psi(B_r)) &\leq \frac{1}{2} \int_{\Psi(B_r)} |DL|^2 + Q \int_{\Psi(B_r)} \langle \boldsymbol{\eta} \circ L, D_{\xi_1} \xi_1 + D_{\xi_2} \xi_2 \rangle \\ &\leq \frac{1}{2} \int_{B_r} |D\mathcal{L}|^2 + C\text{Err}_2(\mathcal{L}, B_r). \end{aligned} \quad (3.13)$$

Combining (3.12) and (3.13) we conclude (3.5).

3.2. Proof of Proposition 3.2 case (b): T semicalibrated. We proceed as in the previous step and define the current Z as in (3.6). If S is any current such that

$$\partial S = T - Z = \mathbf{T}_{\mathcal{F}|_{B_r}} - \mathbf{T}_G = \mathbf{T}_{F|\Psi(B_r)} - \mathbf{T}_G,$$

then the semicalibrated condition gives

$$\mathbf{M}(T) \leq \mathbf{M}(Z) + S(d\omega),$$

where ω is the calibrating form. In particular, in order to conclude the proof it suffices to find an S such that

$$|S(d\omega)| \leq C\text{Err}_1(\mathcal{N}, B_r) + C\text{Err}_2(\mathcal{L}, B_r) + \frac{C}{r} \int_{B_r} |\mathcal{L}|^2 : \quad (3.14)$$

combining the latter inequality with the estimates of the previous subsection we reach the desired inequality.

We first define $H_i : [0, 1] \times \Psi(B_r) \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+n})$ for $i = 1, 2$ by

$$\begin{aligned} [0, 1] \times \Psi(B_r) \ni (t, p) &\mapsto H_1(t, p) := \sum_{i=1}^Q \llbracket p + t N_i(p) \rrbracket \in \mathcal{A}_Q(\mathbb{R}^{2+n}) \\ [0, 1] \times \Psi(B_r) \ni (t, p) &\mapsto H_2(t, p) := \sum_{i=1}^Q \llbracket p + (1-t) L_i(p) \rrbracket \in \mathcal{A}_Q(\mathbb{R}^{2+n}). \end{aligned}$$

We choose $S := S_1 + S_2$, where $S_i := \mathbf{T}_{H_i}$ for $i = 1, 2$. Thanks to the homotopy formula in [5], we get

$$\begin{aligned} \partial S_1 &= \mathbf{T}_{F|\Psi(B_r)} - Q \llbracket \mathcal{M} \rrbracket - \mathbf{T}_{H_1|_{[0,1] \times \Psi(\partial B_r)}}, \\ \partial S_2 &= Q \llbracket \mathcal{M} \rrbracket - \mathbf{T}_{G|\Psi(B_r)} + \mathbf{T}_{H_2|_{[0,1] \times \Psi(\partial B_r)}}. \end{aligned}$$

On the other hand since $N = L$ on $\Psi(\partial B_r)$, we conclude $\partial S = \partial(S_1 + S_2) = T - Z$.

We next estimate $|S_1(d\omega)|$ and $|S_2(d\omega)|$. Since the estimates are analogous, we give the details only for the first. We start from the formula

$$S_1(d\omega) = \int_{\Psi(B_r)} \int_0^1 \sum_{i=1}^Q \langle \vec{\zeta}_i(t, p), d\omega((H_1)_i(t, p)) \rangle d\mathcal{H}^2(p) dt,$$

with

$$\begin{aligned} \vec{\zeta}_i(t, p) &= (\xi_1 + t \nabla_{\xi_1} N_i(p)) \wedge (\xi_2 + t \nabla_{\xi_2} N_i(p)) \wedge N_i(p) \\ &=: \xi_1 \wedge \xi_2 \wedge N_i(p) + \vec{E}_i(t, p), \end{aligned}$$

and

$$|\vec{E}_i(t, p)| \leq C (|DN|(p) + |DN|^2(p)) |N|(p). \quad (3.15)$$

Next we note that

$$d\omega((H_1)_i(t, p)) = d\omega(p) + I(t, p), \quad (3.16)$$

where $I(t, p)$ can be estimated by

$$|I(t, p)| = |d\omega((H_1)_i(t, p)) - d\omega(p)| \leq C \|D^2\omega\|_{L^\infty} |N|(p). \quad (3.17)$$

Therefore, we have

$$\begin{aligned} \left| \sum_{i=1}^Q \langle \vec{\zeta}_i(t, p), d\omega((H_1)_i(t, p)) \rangle \right| &\leq \sum_{i=1}^Q \langle \xi_1 \wedge \xi_2 \wedge N_i(p), d\omega(p) \rangle + \|d\omega\|_{L^\infty} \sum_{i=1}^Q |\vec{E}_i(t, p)| \\ &\quad + C \sum_{i=1}^Q \left((|N_i| + |\vec{E}_i|) |I| \right)(t, p) \\ &\leq C \mathbf{m}_0^{1/2} |\boldsymbol{\eta} \circ N| + C |N|^2(p) + C |DN|(p) |N|(p) + Cr |DN|^2(p), \end{aligned}$$

where we have only used the bound $|N|(p) \leq Cr$ on $\Psi(B_r)$. Arguing similarly for S_2 (observe that we have the bound $|L|(p) \leq Cr$) and estimating $|N||DN| + |L||DL| \leq r^{-1}(|N|^2 + |L|^2) + Cr(|DN|^2 + |DL|^2)$, we achieve

$$\begin{aligned} |S_1(d\omega)| + |S_2(d\omega)| &\leq C \mathbf{m}_0^{1/2} \int_{\Psi(B_r)} (|\boldsymbol{\eta} \circ N| + |\boldsymbol{\eta} \circ L|) + Cr^{-1} \int_{\Psi(B_r)} (|N|^2 + |L|^2) \\ &\quad + Cr \int_{\Psi(B_r)} (|DN|^2 + |DL|^2), \end{aligned}$$

and we conclude as above by a change of variable and Theorem 2.6.

3.3. Proof of Proposition 3.2 in case (c): T is the cross-section of a three dimensional area minimizing cone. Recall that in this case $\text{spt}(T) \subset \partial \mathbf{B}_R(p_0)$, where $p_0 = (0, \dots, 0, R) = Re_{n+2}$ and $R^{-1} \leq \mathbf{m}_0^{1/2}$. For the computations of this subsection it is indeed convenient to change coordinates so that p_0 is in fact the origin, whereas $\Psi(0, 0)$ is the point $(0, \dots, 0, -R)$. In these new coordinates we then have $\mathcal{M}, \text{spt}(T), \text{Im}(\mathcal{F}) \subset \partial \mathbf{B}_R(0)$. These coordinates will however be used only in here, whereas in the next sections we will return to the usual ones.

We introduce the following notation: $\mathcal{C}(r)$ is the cone over $\Psi(B_r)$ with vertex 0, i.e.

$$\mathcal{C}(r) := \{\rho p \in \mathbb{R}^{n+2} : \rho \in [0, 1], p \in \Psi(B_r)\},$$

with the orientation compatible with that of $0 \llcorner \llbracket \mathcal{M} \rrbracket$. We extend F to $\tilde{F} : \mathcal{C}(r) \rightarrow \mathcal{A}_Q(\mathbb{R}^{n+2})$ by setting $\tilde{F}(\rho p) := \rho F(p)$ for every $p \in \Psi(B_r)$.

In order to estimate the Dirichlet energy of N in terms of that of L , we construct a suitable function $K : \mathcal{C}(r) \rightarrow \mathcal{A}_Q(\mathbb{R}^{n+2})$ (depending on L and N) such that $K|_{\partial \mathcal{C}(r)} = \tilde{F}|_{\partial \mathcal{C}(r)}$: we can then test the minimizing property of $0 \llcorner T$ comparing its mass with that of the current

$$Z := 0 \llcorner T - \mathbf{T}_{\tilde{F}} + \mathbf{T}_K = 0 \llcorner (T - \mathbf{T}_{F|_{\Psi(B_r)}}) + \mathbf{T}_K$$

which is easily recognized to satisfy $\partial Z = \partial(0 \llcorner T)$. In particular, using the minimality of $0 \llcorner T$, we conclude

$$R^{-1} \mathbf{M}(0 \llcorner \mathbf{T}_{F|_{\Psi(B_r)}}) \leq R^{-1} \mathbf{M}(\mathbf{T}_K) + C \text{Err}_1(\mathcal{N}, B_r). \quad (3.18)$$

We consider the space of parameters $[0, 1] \times B_r$ and recall that the points in \mathfrak{B}_Q are identified by two complex coordinates $(z, w) \in \mathbb{C} \times \mathbb{C}$. For the definition of K we need to introduce the following sets

$$A_1 := \left\{ (\rho, z, w) \in [0, 1] \times B_r : 1 - r \leq \rho \leq 1, |z| \leq \frac{\rho + 2r - 1}{2} \right\}, \quad (3.19)$$

$$A_2 := \left\{ (\rho, z, w) \in [0, 1] \times B_r : 1 - 2r \leq \rho \leq 1 - r, |z| \leq \frac{1 - \rho}{2} \right\}, \quad (3.20)$$

$$B := [1 - 2r, 1] \times B_r \setminus (A_1 \cup A_2), \quad (3.21)$$

We then define the function $\mathcal{H} : [0, 1] \times B_r \rightarrow \mathcal{A}_Q(\mathbb{R}^{n+2})$ given by

$$\mathcal{H}(\rho, z, w) := \begin{cases} \rho \mathcal{L}(z, w) & \text{if } \rho \leq 1 - 2r, \\ \rho l_1(\rho) \mathcal{N}\left(\frac{2rz}{\rho+2r-1}, \frac{(2r)^{1/Q}}{(\rho+2r-1)^{1/Q}} w\right) & \text{if } (\rho, z, w) \in A_1, \\ -\rho l_1(\rho) \mathcal{L}\left(\frac{2rz}{1-\rho}, \frac{(2r)^{1/Q}}{(1-\rho)^{1/Q}} w\right) & \text{if } (\rho, z, w) \in A_2, \\ \rho l_2(|z|) \mathcal{N}\left(\frac{rz}{|z|}, \frac{r^{1/Q}}{|z|^{1/Q}} w\right) & \text{if } (\rho, z, w) \in B, \end{cases} \quad (3.22)$$

where $l_1, l_2 : \mathbb{R} \rightarrow \mathbb{R}$ are the affine functions

$$l_1(t) := \frac{t+r-1}{r} \quad \text{and} \quad l_2(t) := \frac{2t-r}{r}. \quad (3.23)$$

The following are simple properties of \mathcal{H} which can be easily verified:

- (1) $\mathcal{H}(1, z, w) = \mathcal{N}(z, w)$ for every $(z, w) \in B_r$, as $(1, z, w) \in A_1$ and $l_1(1) = 1$;
- (2) $\mathcal{H}(\rho, z, w) = \rho \mathcal{N}(z, w)$ for every $\rho \in [0, 1]$ and for every z with $|z| = r$, as $\mathcal{L}|_{\partial B_r} = \mathcal{N}|_{\partial B_r}$ and $l_2(r) = 1$;
- (3) \mathcal{H} is well-defined and continuous, as $\mathcal{H} \equiv 0$ in $A_1 \cap A_2$ from $l_1(1-r) = 0$,

$$\mathcal{H}(\rho, z, w) = \rho \frac{\rho+r-1}{r} \mathcal{N}\left(\frac{rz}{|z|}, \frac{r^{1/Q}}{|z|^{1/Q}} z\right) \quad \text{in } A_1 \cap \partial B,$$

and

$$\mathcal{H}(\rho, z, w) = \rho \frac{\rho+r-1}{r} \mathcal{N}\left(\frac{rz}{|z|}, \frac{r^{1/Q}}{|z|^{1/Q}} w\right) \quad \text{in } A_2 \cap \partial B.$$

The competitor map $K : \mathcal{C}(r) \rightarrow \mathcal{A}_Q(\mathbb{R}^{n+2})$ is now given by

$$K(\rho p) := \sum_{i=1}^Q \llbracket \rho p + H_i(\rho p) \rrbracket \quad \text{with } H(\rho p) := \mathcal{H}(\rho, \Psi^{-1}(p)).$$

Note that by (1) and (2) above it follows that $K|_{\partial \mathcal{C}(r)} = \tilde{F}|_{\partial \mathcal{C}(r)}$.

We start now estimating the masses of the various currents introduced above. Since $\text{spt}(\mathbf{T}_F) \subset \partial \mathbf{B}_R(0)$, it follows that $\mathbf{M}(0 \llcorner \mathbf{T}_F) = R \mathbf{M}(\mathbf{T}_F)/3$ and, by the expansion of the mass of \mathbf{T}_F , we have that

$$\mathbf{M}(\mathbf{T}_F|_{\Psi(B_r)}) \geq Q \mathcal{H}^2(\Psi(B_r)) + \frac{1}{2} \int_{B_r} |D\mathcal{N}|^2 - C \text{Err}_1(\mathcal{N}, B_r). \quad (3.24)$$

Combining the latter estimate with (3.18) we conclude

$$\int_{B_r} |D\mathcal{N}|^2 \leq 6R^{-1} \mathbf{M}(\mathbf{T}_K) - 2Q \mathcal{H}^2(\Psi(B_r)) + C \text{Err}_1(\mathcal{N}, B_r). \quad (3.25)$$

For what concerns the mass of \mathbf{T}_K , recalling that $p + \text{spt}(L(p)) \in \partial \mathbf{B}_R(0)$ for every $p \in \Psi(B_r)$, we deduce that

$$\mathbf{M}(\mathbf{T}_K \llcorner \mathbf{B}_{R(1-2r)}) = \mathbf{M}(0 \llcorner \mathbf{T}_G \llcorner \mathbf{B}_{R(1-2r)}) = R \frac{(1-2r)^3 \mathbf{M}(\mathbf{T}_G)}{3}$$

and

$$\mathbf{M}(\mathbf{T}_G) \leq Q \mathcal{H}^2(\Psi(B_r)) + \frac{1}{2} \int_{B_r} |D\mathcal{L}|^2 + \text{Err}_2(\mathcal{L}, B_r).$$

In particular we conclude

$$6R^{-1}\mathbf{M}(\mathbf{T}_K \llcorner \mathbf{B}_{R(1-2r)}) \leq 2Q(1-2r)^3 \mathcal{H}^2(\Psi(B_r)) + \int_{B_r} |D\mathcal{L}|^2 + \text{Err}_2(\mathcal{L}, B_r). \quad (3.26)$$

Next we pass to estimating $\mathbf{M}(\mathbf{T}_K \llcorner \mathbf{B}_R \setminus \mathbf{B}_{R(1-2r)})$. In order to carry on our estimates we use the area formula for multifunctions, cf. [5]. In particular we fix an orthonormal frame ξ_1, ξ_2 for \mathcal{M} as in the proof of case (a) and we let $\xi_3 = R^{-1}\partial_t$ be normal to them in $T\mathcal{C}(r)$, i.e. pointing in the radial direction of the cone. We then have

$$\mathbf{M}(\mathbf{T}_K \llcorner (\mathbf{B}_R \setminus \mathbf{B}_{R(1-2r)})) = \int_{\mathcal{C}(r)} \sum_i \underbrace{|(\xi_1 + DH_i \cdot \xi_1) \wedge (\xi_2 + DH_i \cdot \xi_2) \wedge (\xi_3 + DH_i \cdot \xi_3)|}_{(A)}.$$

Using the Taylor expansion for (A), cf. [5], we can bound

$$\begin{aligned} R^{-1}\mathbf{M}(\mathbf{T}_K \llcorner (\mathbf{B}_R \setminus \mathbf{B}_{R(1-2r)})) &\leq QR^{-1} \mathcal{H}^3(\mathcal{C}(r) \cap \mathbf{B}_1 \setminus \mathbf{B}_{1-2r}) \\ &+ QR^{-1} \int_{1-2r}^1 \int_{\Psi(B_r)} \frac{d}{dt} [(\boldsymbol{\eta} \circ H)(tp)] t^2 dt \\ &+ QR^{-1} \int_{1-2r}^1 \int_{\Psi(B_r)} \sum_{i=1}^2 \langle \nabla_{\xi_i}(\boldsymbol{\eta} \circ H), \xi_i \rangle t^2 dt + CR^{-1} \int_{1-2r}^1 \int_{\Psi(B_r)} |DH|^2 t^2 dt. \end{aligned} \quad (3.27)$$

The linear terms can be integrated by parts: since $\nabla_p(\boldsymbol{\eta} \circ H)(tp) = \frac{d}{dt}(\boldsymbol{\eta} \circ H)(tp)$, we have

$$\begin{aligned} \int_{1-2r}^1 \int_{\Psi(B_r)} \frac{d}{dt} [(\boldsymbol{\eta} \circ H)(tp)] t^2 dt &= \int_{\Psi(B_r)} \langle (\boldsymbol{\eta} \circ H)(p) - (1-2r)^2(\boldsymbol{\eta} \circ H)((1-2r)p), p \rangle \\ &- 2 \int_{1-2r}^1 \int_{\Psi(B_r)} \langle (\boldsymbol{\eta} \circ H)(tp), p \rangle t dt \end{aligned} \quad (3.28)$$

$$\int_{1-2r}^1 \int_{\Psi(B_r)} \sum_{i=1}^2 \langle \nabla_{\xi_i}(\boldsymbol{\eta} \circ H), \xi_i \rangle t^2 dt = - \int_{1-2r}^1 \int_{\Psi(B_r)} \langle (\boldsymbol{\eta} \circ H), H_{\mathcal{M}} \rangle t^2 dt. \quad (3.29)$$

Therefore, by a simple change of coordinates we can estimate

$$R^{-1}\mathbf{M}(\mathbf{T}_K \llcorner (\mathbf{B}_R \setminus \mathbf{B}_{R(1-r)})) \leq \frac{Q(1-(1-2r)^3)}{3} \mathcal{H}^2(\Psi(B_r)) \quad (3.30)$$

$$+ C\mathbf{m}_0^{1/2} \int_{B_r} (|\boldsymbol{\eta} \circ \mathcal{N}| + |\boldsymbol{\eta} \circ \mathcal{L}|) + C\mathbf{m}_0^{1/2} \int_{1-2r}^1 \int_{B_r} |D\mathcal{H}|^2(t, z, w) dz dt \quad (3.31)$$

$$+ C\mathbf{m}_0^{1/2} \int_{1-2r}^1 \int_{B_r} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{H}|(t, z, w) dz dt. \quad (3.32)$$

In order to bound the various integrands of (3.30), we start with the following general remark. Assume that $\chi : [1 - 2r, 1] \times B_r \rightarrow [0, +\infty)$ has the structure

$$\chi(\rho, x, y) = \begin{cases} \chi_1 \left(\frac{2rz}{\rho+2r-1}, \frac{(2r)^{1/Q}}{(\rho+2r-1)^{1/Q}} w \right) & \text{if } (\rho, z, w) \in A_1, \\ \chi_2 \left(\frac{2rz}{1-\rho}, \frac{(2r)^{1/Q}}{(1-\rho)^{1/Q}} w \right) & \text{if } (\rho, z, w) \in A_2, \\ \chi_3 \left(\frac{rz}{|z|}, \frac{r^{1/Q}}{|z|^{1/Q}} w \right) & \text{if } (\rho, z, w) \in B, \end{cases} \quad (3.33)$$

for some $\chi_1, \chi_2, \chi_3 : B_r \rightarrow [0, +\infty)$. Then one can compute the integral of χ in the following way:

$$\int_{1-2r}^1 \int_{B_r} \chi(t, z, w) dz dt = \int_{A_1} \chi(t, z, w) dz dt + \int_{A_2} \chi(t, z, w) dz dt + \int_B \chi(t, z, w) dz dt,$$

and one can easily compute that

$$\begin{aligned} \int_{A_1} \chi(t, z, w) dz dt &= \int_{1-r}^1 \int_{B_{\frac{t+2r-1}{2}}} \chi_1(t, z, w) dz dt \\ &= \int_{1-r}^1 \int_{B_{\frac{t+2r-1}{2}}} \chi_1 \left(\frac{2rz}{t+2r-1}, \frac{(2r)^{1/Q}}{(t+2r-1)^{1/Q}} w \right) dz dt \\ &= \int_{1-r}^1 \left(\frac{t+2r-1}{2r} \right)^2 \int_{B_r} \chi_1(z, w) dz dt \leq r \int_{B_r} \chi_1(z, w) dz dt. \end{aligned} \quad (3.34)$$

Similarly

$$\int_{A_2} \chi(t, z, w) dz dt \leq r \int_{B_r} \chi_2(z, w) dz dt, \quad (3.35)$$

and

$$\begin{aligned} \int_B \chi(t, z, w) dz dt &= \int_{1-r}^1 dt \int_{\frac{t+2r-1}{2}}^r \frac{s}{r} ds \int_{\partial B_r} \chi_3(z, w) dz \\ &\quad + \int_{1-2r}^{1-r} \int_{\frac{1-t}{2}}^r \frac{s}{r} ds \int_{\partial B_r} \chi_3(z, w) dz \leq r^2 \int_{\partial B_r} \chi_3(z, w) dz. \end{aligned} \quad (3.36)$$

By direct computations one verifies that the integrands in (3.30) are all bounded from above by functions χ with the structure (3.33): in particular,

(i) $|z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{H}|(t, z, w) \leq \chi(t, z, w)$ if we choose

$$\chi_1(z, w) = \chi_3(z, w) = |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{X}|(z, w) \quad \text{and} \quad \chi_2(z, w) = |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{L}|(z, w);$$

(ii) $|D\mathcal{H}|^2(t, z, w) \leq \chi(t, z, w)$ if we choose

$$\begin{aligned} \chi_1(z, w) = \chi_3(z, w) &= \frac{C}{r^2} |\mathcal{X}|^2(z, w) + C |D\mathcal{X}|^2(z, w) \\ \chi_2(z, w) &= \frac{C}{r^2} |\mathcal{L}|^2(z, w) + C |DL|^2(z, w). \end{aligned}$$

for some dimensional constant $C > 0$.

It then turns out from (3.34), (3.35), (3.36) and (i), (ii), (iii) that

$$6R^{-1}\mathbf{M}(\mathbf{T}_K \llcorner (\mathbf{B}_R \setminus \mathbf{B}_{R(1-r)})) \leq Q(1 - (1 - 2r)^3) \mathcal{H}^2(\Psi(B_r)) + C \text{Err}_1(\mathcal{N}, B_r) + C \text{Err}_2(\mathcal{L}, B_r). \quad (3.37)$$

Summing (3.37) and (3.26) we conclude

$$6R^{-1}\mathbf{M}(\mathbf{T}_K) \leq 2Q\mathcal{H}^2(\Psi(B_r)) + \int_{B_r} |D\mathcal{L}|^2 + C \text{Err}_1(\mathcal{N}, B_r) + C \text{Err}_2(\mathcal{L}, B_r).$$

Combining the latter estimate with (3.25) we conclude the proof.

4. HARMONIC COMPETITOR

The most natural choice for the competitor \mathcal{L} is a suitable “harmonic” extension of the boundary value $\mathcal{N}|_{\partial B_r}$. Following the ideas of [3] we estimate carefully the energy of such competitor. To this purpose it is useful to introduce “polar” coordinates with center 0 in \mathfrak{B} and split accordingly the Dirichlet integrand in radial and angular parts. More precisely, consider $(z_0, w_0) = ((\xi_0, \zeta_0), w_0) \in \partial B_r$ and take, locally, the standard flat coordinates $z = (x_1, x_2)$ of Definition 1.3. We then denote by ν the exterior unit vector normal to ∂B_r at (z_0, w_0) and by τ the corresponding tangent unit vector obtained by rotating ν of an angle $\pi/2$ in the counterclockwise direction, namely

$$\nu := |z_0|^{-1} \left(\xi_0 \frac{\partial}{\partial x_1} + \zeta_0 \frac{\partial}{\partial x_2} \right) \quad \text{and} \quad \tau := |z_0|^{-1} \left(-\zeta_0 \frac{\partial}{\partial x_1} + \xi_0 \frac{\partial}{\partial x_2} \right).$$

The directional derivatives of any (multi)function f on \mathfrak{B} gives then two (multi)functions

$$D_\nu f = \sum_i \llbracket Df_i \cdot \nu \rrbracket \quad \text{and} \quad D_\tau f = \sum_i \llbracket Df_i \cdot \tau \rrbracket.$$

The Dirichlet integrand $|Df|^2$ enjoys then the splitting

$$|Df|^2 = |D_\nu f|^2 + |D_\tau f|^2.$$

For the rigorous justification of these identities see [4].

Proposition 4.1. *There are constants $C > 0$, $\sigma > 0$ such that, for every $r \in (0, 1)$ there exists a competitor $\mathcal{L}: B_r \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+n})$ for \mathcal{N} with the following additional properties:*

- (i) $\text{Lip}(\mathcal{L}) \leq C_{3.2}^{-1}$, $\|\mathcal{L}\|_0 \leq Cr$.
- (ii) *The following estimates hold:*

$$\int_{B_r} |D\bar{\mathcal{L}}|^2 \leq Cr \int_{\partial B_r} |D\bar{\mathcal{N}}|^2 \leq Cr \mathbf{D}'(r), \quad (4.1)$$

$$\int_{B_r} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{L}| \leq Cr^{\gamma_0} \int_{\partial B_r} |\boldsymbol{\eta} \circ \mathcal{N}| + C \mathbf{H}(r). \quad (4.2)$$

(iii) For every $a > 0$ there exists $b_0 > 0$ such that, for all $b \in (0, b_0)$, the following estimate holds:

$$(2a + b) \int_{B_r} |D\bar{\mathcal{L}}|^2 \leq r \int_{\partial B_r} |D_\tau \mathcal{X}|^2 + \frac{a(a+b)}{r} \int_{\partial B_r} |\mathcal{X}|^2 + Cr^{1+\sigma} \mathbf{D}'(r). \quad (4.3)$$

Using this competitor in Proposition 3.2, we then infer the following corollary.

Corollary 4.2. For every $r \in (0, 1)$ the following inequality holds

$$\mathbf{D}(r) \leq Cr \mathbf{D}'(r) + C \mathbf{H}(r) + C \mathbf{F}(r) + C \mathbf{m}_0^{1/2} r^{\gamma_0} \int_{\partial B_r} |\boldsymbol{\eta} \circ \mathcal{X}|. \quad (4.4)$$

For every $a > 0$ there exists $b_0 > 0$ such that, for all $b \in (0, b_0)$ and all $r \in]0, 1[$

$$\mathbf{D}(r) \leq (1 + Cr) \left[\frac{r}{(2a + b)} \int_{\partial B_r} |D_\tau \mathcal{X}|^2 + \frac{a(a+b)}{r(2a + b)} \mathbf{H}(r) \right] + C \mathcal{E}_{QM}(r) + Cr^{1+\sigma} \mathbf{D}'(r), \quad (4.5)$$

with

$$\mathcal{E}_{QM}(r) \leq \Lambda(r)^{\eta_0} \mathbf{D}(r) + \mathbf{F}(r) + \mathbf{H}(r) + \mathbf{m}_0^{1/2} r^{\gamma_0} \int_{\partial B_r} |\boldsymbol{\eta} \circ \mathcal{X}|.$$

Proof of Corollary 4.2. Recalling that $\mathbf{H}(r) \leq Cr \|\mathcal{X}\|_{\partial B_r}^2 \leq Cr^{3+\gamma_0}$ we easily infer that $\Lambda(r) \leq Cr^2$ and thus the inequalities follow readily from Proposition 3.2 and Proposition 4.1. \square

4.1. Proof of Proposition 4.1: Step 1. First of all we observe that it suffices to exhibit $\bar{\mathcal{L}}$, as \mathcal{L} can be recovered from it via the formula (3.1). Moreover, it suffices to show the estimates with $\bar{\mathcal{X}}$ in place of \mathcal{X} in the right hand side, because we obviously have $|\bar{\mathcal{X}}| \leq |\mathcal{X}|$ and $|D\bar{\mathcal{X}}| \leq |D\mathcal{X}|$. Next we wish to relate $\boldsymbol{\eta} \circ \mathcal{L}$ and $\boldsymbol{\eta} \circ \bar{\mathcal{L}}$ for two maps satisfying the relation (3.1). Note that by a simple Taylor expansion we have

$$|\boldsymbol{\eta} \circ \mathcal{L}| \leq C |\boldsymbol{\eta} \circ \bar{\mathcal{L}}| + C \mathcal{G}(\bar{\mathcal{L}}, \boldsymbol{\eta} \circ \bar{\mathcal{L}})^2,$$

where the constant C depends on the C^2 norm of Ψ_0 . In particular we record the following conclusion:

$$\int_{B_r} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{L}| \leq C \int_{B_r} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \bar{\mathcal{L}}| + C \int_{B_r} |z|^{\gamma_0-1} |\bar{\mathcal{L}}|^2. \quad (4.6)$$

In this step we exhibit an ‘‘harmonic’’¹ competitor \mathcal{H} which satisfies all the requirements of the proposition except for the Lipschitz estimate. In fact we will show that there is a

¹We remark that the competitor used here does not coincide, in general, with the Dirichlet minimizer with boundary value $\bar{\mathcal{X}}|_{\partial B_r}$.

$W^{1,2}$ map $\mathcal{H} : B_r \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+\bar{n}})$ such that

$$\mathcal{H}|_{\partial B_r} = \bar{\mathcal{X}}|_{\partial B_r} \quad \text{and} \quad \|\mathcal{H}\|_{L^\infty(B_r)} \leq Q \|\bar{\mathcal{X}}\|_{L^\infty(\partial B_r)} \quad (4.7)$$

$$\int_{B_r} |D\mathcal{H}|^2 \leq Cr \int_{\partial B_r} |D\bar{\mathcal{X}}|^2 \quad (4.8)$$

$$\int_{B_r} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{H}| \leq Cr^{\gamma_0} \int_{\partial B_r} |\boldsymbol{\eta} \circ \bar{\mathcal{X}}| \quad (4.9)$$

$$\int_{B_r} |z|^{\gamma_0-1} |\mathcal{H}|^2 \leq Cr^{\gamma_0} \int_{\partial B_r} |\bar{\mathcal{X}}|^2 \quad (4.10)$$

$$(2a+b) \int_{B_r} |D\bar{\mathcal{H}}|^2 \leq r \int_{\partial B_r} |D_\tau \bar{\mathcal{X}}|^2 + \frac{a(a+b)}{r} \int_{\partial B_r} |\bar{\mathcal{X}}|^2. \quad (4.11)$$

In these estimates we do not use any of the particular properties of $\bar{\mathcal{X}}$ and indeed for any Lipschitz multivalued map $\bar{\mathcal{X}} : B_r \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+\bar{n}})$ there is such a ‘‘harmonic’’ competitor. Therefore, given the scaling invariance of the estimates, we will assume without loss of generality that $r = 1$.

Let $D_r := \{|z| < r\}$ denote the disk of radius r in \mathbb{R}^2 , which we identify with the complex plane. We start by defining the ‘‘winding map’’ $\mathbf{W} : D_1 \rightarrow \mathfrak{B}$ given (in complex notation) by

$$\mathbf{W}(z) := (z^{\bar{Q}}, z).$$

We then consider the multivalued map $\mathcal{U} := \bar{\mathcal{X}} \circ \mathbf{W}$. Let $\theta \mapsto u(\theta)$ be its trace on $\partial D_1(0)$, which we parametrize with the angle $\theta \in [0, 2\pi]$. According to [4, Proposition 1.5] we can decompose u in a superposition of simple functions $u(\theta) = \sum_{j=1}^J u_j(\theta)$ such that, for every $j = 1, \dots, J$,

$$u_j(\theta) = \sum_{i=1}^{Q_j} \left[\gamma_j \left(\frac{\theta + 2\pi i}{Q_j} \right) \right],$$

where the $\gamma_j : [0, 2\pi] \rightarrow \mathbb{R}^{2+\bar{n}}$ are periodic Lipschitz functions. Next consider the Fourier’s expansion of each γ_j

$$\gamma_j(\theta) = \frac{a_{j,0}}{2} + \sum_{l=1}^{\infty} (a_{j,l} \cos(l\theta) + b_{j,l} \sin(l\theta)),$$

and its harmonic extension, which in polar coordinates (ρ, θ) reads as

$$\zeta_j(\rho, \theta) := \frac{a_{j,0}}{2} + \sum_{l=1}^{\infty} \rho^l (a_{j,l} \cos(l\theta) + b_{j,l} \sin(l\theta)). \quad (4.12)$$

We then can define the ‘‘harmonic’’ competitor for \mathcal{U} , which is the Q -valued map

$$\mathcal{V}(\rho, \theta) := \sum_{j=1}^J \sum_{i=1}^{Q_j} \left[\zeta_j \left(\rho^{1/Q_j}, \frac{\theta + 2\pi i}{Q_j} \right) \right]$$

and the “harmonic” competitor for $\bar{\mathcal{X}}$, which is $\mathcal{H} = \mathcal{V} \circ \mathbf{W}^{-1}$. Observe that the first claim in (4.7) is obvious, whereas the second claim follows from the maximum principle for classical harmonic functions.

Simple computations and the conformality of \mathbf{W} , see for instance [4, Proof of Proposition 5.2], yield

$$\int_{B_1} |D\mathcal{H}|^2 = \int_{D_1} |D\mathcal{V}|^2 = \pi \sum_{j=1}^J \sum_{l=1}^{\infty} l(|a_{j,l}|^2 + |b_{j,l}|^2), \quad (4.13)$$

$$\int_{\partial B_1} |D_\tau \mathcal{H}|^2 = \frac{\pi}{\bar{Q}} \sum_{j=1}^J \sum_{l=1}^{\infty} \frac{l^2}{Q_j} (|a_{j,l}|^2 + |b_{j,l}|^2), \quad (4.14)$$

$$\int_{\partial B_1} |\mathcal{H}|^2 = \pi \bar{Q} \sum_{j=1}^J Q_j \left(\frac{|a_{j,0}|^2}{2} + \sum_{l=1}^{\infty} (|a_{j,l}|^2 + |b_{j,l}|^2) \right). \quad (4.15)$$

Clearly, (4.8) follows from the first and second inequality, with the constant $C = \bar{Q}Q_1 \leq \bar{Q}Q$, assuming that $Q_1 = \max\{Q_1, \dots, Q_j\}$. (4.11) follows from the fact that, for any chosen $a > 0$, if b_0 is sufficiently small and $0 < b < b_0$, then

$$(2a + b)\ell \leq \frac{\ell^2}{\bar{Q}Q_j} + \bar{Q}Q_j \ell a(a + b) \quad \forall \ell \in \mathbb{N}.$$

The latter claim is elementary and the reader can consult, for instance, Step 2 in the proof of [4, Proposition 5.2].

Observe next that $\boldsymbol{\eta} \circ \mathcal{V}$ is the classical harmonic extension of the single-valued function $\boldsymbol{\eta} \circ \mathcal{U}|_{\partial D_1}$. We then have the classical estimates

$$\|\boldsymbol{\eta} \circ \mathcal{V}\|_{L^\infty(D_{2^{1/\bar{Q}}})} + \|\boldsymbol{\eta} \circ \mathcal{V}\|_{L^1(D_1)} \leq C \|\boldsymbol{\eta} \circ \mathcal{U}\|_{L^1(\partial D_1)}.$$

In particular we conclude easily

$$\|\boldsymbol{\eta} \circ \mathcal{H}\|_{L^\infty(B_{1/2})} + \|\boldsymbol{\eta} \circ \mathcal{H}\|_{L^1(B_1 \setminus B_{1/2})} \leq C \int_{\partial B_1} |\boldsymbol{\eta} \circ \bar{\mathcal{X}}|,$$

because the change of variables \mathbf{W}^{-1} is smooth on $B_1 \setminus B_{1/2}$. The integrability of $|z|^{\gamma_0-1}$ on B_1 gives then

$$\int_{B_1} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{H}(z, w)| dz \leq C \|\boldsymbol{\eta} \circ \mathcal{H}\|_{L^\infty(B_{1/2})} + C \|\boldsymbol{\eta} \circ \mathcal{H}\|_{L^1(B_1 \setminus B_{1/2})},$$

which in turn completes the proof of (4.9).

A similar argument proves (4.10). Using the classical theory of single valued harmonic functions we see indeed that $\|\zeta_j\|_{L^2(B_1)} + \|\zeta_j\|_{L^\infty(B_{1/2})} \leq C \|\gamma_j\|_{L^2(\partial B_1)}$ and thus, using the fact that \mathbf{W} is smooth on $B_1 \setminus B_{1/2}$, we conclude that

$$\|\mathcal{H}\|_{L^\infty(B_{1/2})}^2 + \|\mathcal{H}\|_{L^2(B_1 \setminus B_{1/2})}^2 \leq C \int_{\partial B_1} |\bar{\mathcal{X}}|^2.$$

From this we easily conclude (4.10).

4.2. Proof of Proposition 4.1: Step 2. We keep the notation of the previous paragraphs and assume that $\bar{\mathcal{N}}$ is defined in B_1 , after scaling. The specific scaling that we are using is the one which preserves the Lipschitz constant and is given by

$$\bar{\mathcal{N}}(z, w) \mapsto r^{-1}\bar{\mathcal{N}}(rz, r^{1/\bar{Q}}w)$$

and by abuse of notation we keep the symbols $\bar{\mathcal{N}}$, \bar{L} , etc. for all the rescaled maps.

Under this scaling we then have the estimates $\|\bar{\mathcal{N}}\|_{L^\infty} \leq C\mathbf{m}_0^{1/4}r^{\gamma_0/2}$ and $\text{Lip}(\bar{\mathcal{N}}) \leq \mathbf{\Lambda}(r)^{\eta_0}$ and we want to show that we can modify \mathcal{H} to a competitor \bar{L} with $\text{Lip}(\bar{L}) \leq C_{3.2}^{-1}$, satisfying

$$\bar{L}|_{\partial B_1} = \bar{\mathcal{N}}|_{\partial B_1} \quad \text{and} \quad \|\bar{L}\|_{L^\infty(B_1)} \leq C\|\bar{\mathcal{N}}\|_{L^\infty(\partial B_1)} \quad (4.16)$$

$$\int_{B_1} |D\bar{L}|^2 \leq C(1+r^\sigma) \int_{B_1} |D\mathcal{H}|^2 + C\mathbf{\Lambda}(r)^\sigma \int_{\partial B_1} |D\bar{\mathcal{N}}|^2 \quad (4.17)$$

$$\int_{B_r} |z|^{\gamma_0-1} |\bar{L}|^2 \leq C \int_{\partial B_1} |\bar{\mathcal{N}}|^2 \quad (4.18)$$

$$\int_{B_1} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \bar{L}| \leq C \int_{\partial B_1} |\boldsymbol{\eta} \circ \bar{\mathcal{N}}|. \quad (4.19)$$

Observe that the harmonic functions ζ_j defined in (4.12) are Lipschitz in every ball D_{1-t} for $0 < t < 1$ with an estimate of the form

$$\|D\zeta_j\|_{L^\infty(D_{1-t})} \leq \frac{C}{t} \text{Lip}(\gamma_j) \leq \frac{C}{t} \text{Lip}(\bar{\mathcal{N}}) \leq \frac{C\mathbf{\Lambda}(r)^{\eta_0}}{t}. \quad (4.20)$$

They are not Lipschitz up to the boundary ∂D_1 because the Dirichlet to Neumann map $\gamma_j \rightarrow \frac{\partial \zeta_j}{\rho}(1, \cdot)$ does not map L^∞ into L^∞ . However we have the estimate

$$\|D\zeta_j\|_{L^p(D_1)} \leq C_p \|\gamma_j\|_{W^{1,p}(\partial D_1)} \leq C_p \mathbf{\Lambda}(r)^{\eta_0}$$

for every $p < \infty$. In particular, we can bound

$$\|\zeta_j(1-t, \cdot) - \gamma_j\|_{W^{1,1}(\partial D_1)} \leq C_2 t^{1/2} \mathbf{\Lambda}(r)^{\eta_0},$$

which in turn implies

$$\max |\zeta_j(1-t, \theta) - \gamma_j(\theta)| \leq C_2 t^{1/2} \mathbf{\Lambda}(r)^{\eta_0}. \quad (4.21)$$

Choose $t := \mathbf{\Lambda}(r)^{\eta_0/2}$ and define a new map ξ_j as

$$\xi_j(\rho, \theta) := \begin{cases} \zeta_j(\rho, \theta) & \text{for } \rho \leq 1-t \\ \frac{1-\rho}{t} \zeta_j(1-t, \theta) + \frac{\rho-(1-t)}{t} \gamma_j(\theta) & \text{for } 1-t \leq \rho \leq 1. \end{cases}$$

Now, (4.20) and (4.21) imply that $\|D\zeta_j\| \leq C\mathbf{\Lambda}(r)^{\eta_0/2}$. Moreover we obviously have

$$\begin{aligned} \int_{D_1} |D\xi_j|^2 &\leq \int_{D_1} |D\zeta_j|^2 + C\mathbf{\Lambda}(r)^{\eta_0} \left(\int_{\partial D_{1-t}} |D\zeta_j|^2 + \int_{\partial D_1} |D\gamma_j|^2 \right) \\ &\leq \int_{D_1} |D\zeta_j|^2 + Cr\mathbf{\Lambda}(r)^{\eta_0} \int_{\partial B_1} |D\gamma_j|^2. \end{aligned} \quad (4.22)$$

We can now define two “intermediate” maps

$$\mathcal{V}^0(\rho, \theta) := \sum_{j=1}^J \sum_{i=1}^{Q_j} \left[\xi_j \left(\rho^{1/Q_j}, \frac{\theta + 2\pi i}{Q_j} \right) \right]$$

and $\mathcal{L}^0 := \mathcal{V}^0 \circ \mathbf{W}^{-1}$. It is then immediate to see that \mathcal{L}^0 enjoys the bound $\text{Lip}(\mathcal{L}^0) \leq C\mathbf{\Lambda}(r)^{n_0/2}$ on the domain $B_1 \setminus B_{1/4}$ and that all the estimates (4.16), (4.17) and (4.19). On the other hand the differential $D\mathcal{L}^0$ is singular in the origin and in fact it is rather easy to see that we have the bound

$$|D\mathcal{L}^0(z, w)|^2 \leq C|z|^{2-2/(Q\bar{Q})} \int_{B_1} |D\mathcal{L}^0|^2. \quad (4.23)$$

In order to produce $\bar{\mathcal{L}}$ we need to smooth the singularity of \mathcal{L}^0 at the origin. There are several ways to do this and we present here one possibility. First of all we fix $2 < p < 2Q\bar{Q}/(2Q\bar{Q} - 2)$ and observe that (4.23) yields the estimate

$$\int_{B_{3/4}} |D\mathcal{L}^0(z, w)|^p \leq C \left(\int_{B_1} |D\mathcal{L}^0|^2 \right)^{p/2}. \quad (4.24)$$

Next we define

$$M|D\mathcal{L}^0(z, w)| := \sup_{\rho < 1/4} \frac{1}{\rho^2} \int_{B_\rho(z, w)} |D\mathcal{L}^0(z, w)|$$

and let

$$A := \{(z, w) : M|D\mathcal{L}^0(z, w)| \geq c_0\}$$

where c_0 is a constant to be chosen later. Observe that, given the Lipschitz bound for \mathcal{L}^0 outside the origin, for r sufficiently small the set A is contained in $B_{1/2}$. Arguing as in the proof of [4, Proposition 4.4] we have the Lipschitz estimate $\text{Lip}(\mathcal{L}^0) \leq Cc_0$ on $B_1 \setminus A$, where C is a dimensional constant. We can then use the Lipschitz extension of [4, Theorem 1.7] to extend \mathcal{L}^0 to $\bar{\mathcal{L}}$ on A so that $\text{Lip}(\bar{\mathcal{L}}) \leq Cc_0$. Choosing c_0 accordingly we achieve the desired Lipschitz bound on B_1 . As for (4.16) and (4.18) observe that the extension satisfies

$$\|\bar{\mathcal{L}}\|_{L^\infty(B_{1/2})}^2 \leq C\|\mathcal{H}\|_{L^\infty(B_{3/4})}^2$$

and coincides with \mathcal{L}_0 on $B_1 \setminus B_{1/2}$. As for (4.19), it would suffice to show that $|\boldsymbol{\eta} \circ \bar{\mathcal{L}}| \leq C|\boldsymbol{\eta} \circ \bar{\mathcal{X}}|$. This can be easily achieved in the following way: we make a Lipschitz extension of \mathcal{L}^0 , subtract from each sheet the average and then sum back to each sheet a Lipschitz extension of $\boldsymbol{\eta} \circ \mathcal{L}^0$.

As for (4.17) we compute

$$\begin{aligned} \int |D\bar{\mathcal{L}}|^2 &\leq \int |D\mathcal{L}^0|^2 + Cc_0^2|A| \leq \int |D\mathcal{L}^0|^2 + Cc_0^{2-p} \int_{B_{3/4}} |D\mathcal{L}^0|^p \\ &\leq \int |D\mathcal{L}^0|^2 \left(1 + Cc_0^{2-p} \left(\int |D\mathcal{L}^0|^2 \right)^{p/2-1} \right). \end{aligned} \quad (4.25)$$

Observe that $p/2 - 1 > 0$ and that by (4.22) and (4.8)

$$\int |D\mathcal{L}^0|^2 \leq \int |D\mathcal{H}|^2 + C\Lambda(r)^{\sigma/2} \int_{\partial B_1} |D\bar{\mathcal{N}}|^2 \leq C \int_{\partial B_1} |D\bar{\mathcal{N}}|^2 \leq Cr^\sigma.$$

so that

$$\int_{B_1} |D\bar{\mathcal{L}}|^2 \leq (1 + Cr^\sigma) \int_{B_1} |D\mathcal{H}|^2 + Cr^\sigma \int_{\partial B_1} |D\bar{\mathcal{N}}|^2 \stackrel{(4.8)}{\leq} \int_{B_1} |D\mathcal{H}|^2 + Cr^\sigma \int_{\partial B_1} |D\bar{\mathcal{N}}|^2.$$

5. OUTER VARIATIONS AND THE POINCARÉ INEQUALITY

In this section we begin to exploit the first variations of the area functional on T in conjunction with the estimates of the previous section. The main conclusion will be the following Poincaré inequality:

Theorem 5.1 (Poincaré inequality). *There exists a constant $C_{5.1} > 0$ such that if r is sufficiently small, then*

$$\mathbf{H}(r) \leq C_{5.1} r \mathbf{D}(r). \quad (5.1)$$

We record however the two main tools used to prove Theorem 5.1, since they will be useful in the future. The first one is an elementary computation. In order to state it we introduce the quantity

$$\mathbf{E}(r) := \int_{\partial B_r} \sum_{j=1}^Q \langle \mathcal{N}_j, D_\nu \mathcal{N}_j \rangle. \quad (5.2)$$

Lemma 5.2. *\mathbf{H} is a Lipschitz function and the following identity holds for a.e. $r \in (0, 1)$*

$$\mathbf{H}'(r) = \frac{\mathbf{H}(r)}{r} + 2\mathbf{E}(r). \quad (5.3)$$

The second identity is a consequence of the first variations of T under specific vector fields, which we call “outer variations”: such variations “stretch” the normal bundle of \mathcal{M} suitably and they are defined using the map \mathcal{N} . In the case of semicalibrated currents it is convenient to modify the Dirichlet energy suitably to gain a new quantity which enjoys better estimates. Thus, from now on Ω will denote \mathbf{D} in the cases (a) and (c) of Definition 0.1, whereas in the case (b) it will be given by

$$\begin{aligned} \Omega(r) &:= \mathbf{D}(r) + \mathbf{L}(r) \\ &:= \mathbf{D}(r) + \int_{\Psi(B_r)} \sum_{i=1}^Q \langle \xi_1(p) \wedge D_{\xi_2} N_i(p) \wedge N_i(p) + D_{\xi_1} N_i(p) \wedge \xi_2(p) \wedge N_i(p), d\omega(p) \rangle dp. \end{aligned}$$

Proposition 5.3 (Outer variations). *There exist constants $C_{5.3} > 0$ and $\kappa > 0$ such that, if $r > 0$ is small enough, then the inequality*

$$|\Omega(r) - \mathbf{E}(r)| \leq C_{5.3} \mathcal{E}_{OV}(r) \quad (5.4)$$

holds with

$$\mathcal{E}_{OV}(r) = \mathbf{\Lambda}(r)^\kappa \left(\mathbf{D}(r) + \frac{\mathbf{H}(r)}{r} + r\mathbf{D}'(r) \right) + \mathbf{F}(r) + r^{1+\gamma_0} \frac{d}{dr} \|T - \mathbf{T}_F\|(\mathbf{p}^{-1}(\Psi(B_r))). \quad (5.5)$$

Moreover

$$|\mathbf{L}(r)| \leq C \mathbf{m}_0^{1/2} r^{2-\gamma_0} \mathbf{D}(r) + C \mathbf{m}_0^{1/2} \mathbf{F}(r). \quad (5.6)$$

5.1. Proof of Lemma 5.2. The Lipschitz regularity of \mathbf{H} follows from the Lipschitz regularity of \mathcal{N} . Consider next the map $i_r : \mathfrak{B} \rightarrow \mathfrak{B}$ given by $i_r(z, w) = (rz, r^{1/Q}w)$. By a simple change of variables we compute

$$\mathbf{H}(r) = \int_{\partial B_1} |\mathcal{N}|^2(i_r(z', w')) r.$$

The formula (5.3) is then an elementary computation using the chain rule for multifunctions, cf. [4].

5.2. Proof of Proposition 5.3. The inequality (5.6) is a simple consequence of

$$|\mathbf{L}(r)| \leq C \mathbf{m}_0^{1/2} \int_{B_r} |D\mathcal{N}| |\mathcal{N}| \leq C \mathbf{m}_0^{1/2} \int_{B_r} |z|^{2-\gamma_0} |D\mathcal{N}|^2 + C \mathbf{m}_0^{1/2} \int_{B_r} |z|^{\gamma_0-2} |\mathcal{N}|^2.$$

In order to show (5.4) we fix a test function $\phi \in C_c^\infty(\mathbb{R})$, nonnegative, symmetric, with support in $] -1, 1[$ and monotone decreasing on $[0, 1]$. We then follow [8, Section 3.3] and, having fixed r , we define the vector field X^o on $\mathbf{V}_{u,a}$ via

$$X^o(p) := \varphi(\mathbf{p}(p))(p - \mathbf{p}(p)) \quad \text{where} \quad \varphi(\Psi(z, w)) = \phi\left(\frac{|z|}{r}\right).$$

For r small enough, by (2.13) we can argue as in [8, Section 3.3] and deduce via the change of coordinates given by Ψ , that

$$\delta \mathbf{T}_F(X) = \int_{\mathfrak{B}} \phi\left(\frac{|z|}{r}\right) |D\mathcal{N}|^2 + r^{-1} \int_{\mathfrak{B}} \phi'\left(\frac{|z|}{r}\right) \sum_{j=1}^Q \langle \mathcal{N}_j, D_\nu \mathcal{N}_j \rangle + \sum_{i=1}^3 \text{Err}_i^o, \quad (5.7)$$

with

$$\text{Err}_1^o = \left| \int_{\mathcal{M}} \varphi \langle H_{\mathcal{M}}, \boldsymbol{\eta} \circ N \rangle \right| \leq C \mathbf{m}_0^{1/2} \int_{B_r} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{N}| \stackrel{(2.14)}{\leq} C \mathbf{\Lambda}^{\eta_0}(r) \mathbf{D}(r) + C \mathbf{F}(r), \quad (5.8)$$

$$\text{Err}_2^o \leq C \int_{\mathcal{M}} |\varphi| |A_{\mathcal{M}}|^2 |N|^2 \leq C \mathbf{F}(r), \quad (5.9)$$

$$\begin{aligned}
\text{Err}_3^o &\leq C \int_{\mathcal{M}} \left(|\varphi| (|DN|^2 |N| |A_{\mathcal{M}}| + |DN|^4) + |D\varphi| (|DN|^3 |N| + |DN| |N|^2 |A_{\mathcal{M}}|) \right) \\
&\leq C \int_{B_r} \left[\left(\frac{|\mathcal{N}|^2}{|z|^{2-2\gamma_0}} + |D\mathcal{N}|^4 \right) - r^{-1} \phi' \left(\frac{|z|}{r} \right) r^{1+\gamma_0} |D\mathcal{N}|^3 - r^{-1} \phi' \left(\frac{|z|}{r} \right) |D\mathcal{N}| \frac{|\mathcal{N}|^2}{|z|^{1-\gamma_0}} \right] \\
&\stackrel{(2.13)\&(2.12)}{\leq} C \mathbf{\Lambda}^{\eta_0}(r) \mathbf{D}(r) + C \mathbf{F}(r) - C \mathbf{\Lambda}(r)^{\eta_0} \int_{B_r} r^{-1} \phi' \left(\frac{|z|}{r} \right) \frac{|\mathcal{N}|^2}{|z|^{1-\gamma_0}} \\
&\quad - C r^{1+\gamma_0} \mathbf{\Lambda}^{\eta_0} \int_{B_r} r^{-1} \phi' \left(\frac{|z|}{r} \right) |D\mathcal{N}|^2. \tag{5.10}
\end{aligned}$$

(We recall that $\phi' \leq 0$ on $[0, 1]$).

We next drop the superscript from X^o and we distinguish two situations:

- In the cases (a) and (c) of Definition 0.1, we denote by X^\perp and X^T the projections of X on the normal and the tangential bundle of Σ , respectively. Then $\delta T(X^T) = 0$ and therefore

$$|\delta \mathbf{T}_F(X)| \leq \underbrace{|\delta \mathbf{T}_F(X) - \delta T(X)|}_{\text{Err}_4^o} + \underbrace{|\delta T(X^\perp)|}_{\text{Err}_5^o};$$

- In case (b), since $\delta T(X) = T(dw \lrcorner X)$, we estimate

$$|\delta \mathbf{T}_F(X) - \mathbf{T}_F(dw \lrcorner X)| \leq \underbrace{|\delta \mathbf{T}_F(X) - \delta T(X)| + |T(dw \lrcorner X) - \mathbf{T}_F(dw \lrcorner X)|}_{\text{Err}_4^o}.$$

In both cases we have

$$\begin{aligned}
\text{Err}_4^o &\leq Q \int_{\text{spt}(T) \setminus \text{Im}(F)} |\text{div}_{\bar{T}} X| \, d\|T\| + Q \int_{\text{Im}(F) \setminus \text{spt}(T)} |\text{div}_{\bar{\mathbf{T}}_F} X| \, d\|\mathbf{T}_F\| \\
&\quad + Q \|d\omega\|_\infty \int |X| \, d\|T - \mathbf{T}_F\|,
\end{aligned}$$

where we use the convention that $\omega = 0$ in the cases (a) and (c). We then can estimate

$$\begin{aligned}
\text{Err}_4^o &\leq C \int (\varphi'(\mathbf{p}(p)) |p - \mathbf{p}(p)| + \varphi(\mathbf{p}(p))) \, d\|T - \mathbf{T}_F\| \\
&\stackrel{(2.12)\&(2.15)}{\leq} C \mathbf{\Lambda}^{\eta_0}(r) \mathbf{D}(r) + C \mathbf{F}(r) + C r^{1+\gamma_0} \underbrace{\int |\nabla \varphi(\mathbf{p}(p))| |p - \mathbf{p}(p)| \, d\|T - \mathbf{T}_F\|}_{S(\varphi)}. \tag{5.11}
\end{aligned}$$

In case (b) we have that

$$\mathbf{T}_F(dw \lrcorner X) = \int_{\mathcal{M}} \varphi \sum_{i=1}^Q \langle (\xi_1 + D_{\xi_1} N_i) \wedge (\xi_2 + D_{\xi_2} N_i \cdot \xi_2) \wedge N_i, d\omega(p + N_i(p)) \rangle.$$

Clearly

$$\begin{aligned} & \left| \mathbf{T}_F(d\omega \lrcorner X) - \int_{\mathcal{M}} \varphi \sum_{i=1}^Q \langle (\xi_1 + D_{\xi_1} N_i) \wedge (\xi_2 + D_{\xi_2} N_i \cdot \xi_2) \wedge N_i, d\omega(p) \rangle \right| \\ & \leq C \|d\omega\|_1 \int \varphi |N|^2 \end{aligned}$$

and we can therefore conclude

$$\begin{aligned} & \left| \mathbf{T}_F(d\omega \lrcorner X) - \int_{\mathcal{M}} \varphi \sum_{i=1}^Q \langle \xi_1(p) \wedge D_{\xi_2} N_i(p) \wedge N_i(p) + D_{\xi_1} N_i(p) \wedge \xi_2(p) \wedge N_i(p), d\omega(p) \rangle \right| \\ & \leq C \|d\omega\|_0 \int \varphi |N| |DN|^2 + C \|d\omega\|_0 \int \varphi |\boldsymbol{\eta} \circ N| + C \|d\omega\|_1 \int \varphi |N|^2. \end{aligned}$$

Letting ϕ converge to the characteristic function of the interval $[-1, 1]$, we reach the conclusion (5.4). The only term which needs some care is the term $S(\varphi)$ in (5.11). Note that we can approximate the characteristic function of $[-1, 1]$ with an increasing sequence of functions ϕ_j with the property that $|\phi'_j| \leq Cj$, $0 \leq \phi_j \leq 1$ and $\phi_j \equiv 1$ on $[-1 + 1/j, 1 - 1/j]$. Then we would have

$$\limsup_j S(\varphi_j) \leq C \limsup_j \frac{j}{r} \|T - \mathbf{T}_F\|(\Psi(B_r \setminus B_{r(1-1/j)})) \leq C \frac{d}{dr} \|T - \mathbf{T}_F\|(\Psi(B_r)),$$

by the monotonicity of the function $r \mapsto \|T - \mathbf{T}_F\|(\Psi(B_r))$.

In the cases (a) and (c) we follow the same argument, but we need to bound the additional term Err_5^o . In order to deal with the latter term we argue as in [8, Section 4.1]. In particular we bound

$$\begin{aligned} \text{Err}_5^o & \leq \left| \int \text{div}_{\vec{T}} X^\perp d\|T\| \right| \\ & \leq \underbrace{\int_{\text{spt}(T) \setminus \text{Im}(F)} |\text{div}_{\vec{T}} X| d\|T\| + \int_{\text{Im}(F) \setminus \text{spt}(T)} |\text{div}_{\vec{T}_F} X| d\|\mathbf{T}_F\|}_{I_1} \\ & \quad + \underbrace{\left| \int \langle X^\perp, h(\vec{T}_F(p)) \rangle d\|\mathbf{T}_F\| \right|}_{I_2}, \end{aligned} \tag{5.12}$$

where $h(v_1 \wedge v_2) := \sum_{i=1}^2 A_\Sigma(v_i, v_i)$. Since the projection on the normal to Σ is a C^{2,ε_0} map, X^\perp enjoys the same C^1 bounds as X and I_1 can be controlled as Err_4^o . The term I_2 can be estimated using

$$|X^{\perp}(p)| = \varphi |\mathbf{p}_{T_p \Sigma^\perp}(p - \mathbf{p}(p))| \leq C \mathbf{c}(\Sigma) \varphi |p - \mathbf{p}(p)|^2 \leq C \mathbf{m}_0^{1/2} \varphi |p - \mathbf{p}(p)|^2 \quad \forall p \in \Sigma.$$

In particular we achieve $I_2 \leq C \mathbf{H}(r)$, which concludes the proof.

5.3. Proof of Theorem 5.1. In order to prove the theorem we start estimating the error term \mathbf{F} .

Lemma 5.4. *There exist a constant $C_{5.4} > 0$ (depending on γ_0) such that*

$$\mathbf{F}(r) \leq C_{5.4} r^{\gamma_0-1} \mathbf{H}(r) + C_{5.4} r^{\gamma_0} \mathbf{D}(r) \quad \forall r \in (0, 1). \quad (5.13)$$

Proof. Using (5.3) and an integration by parts we infer that

$$\gamma_0 \int_0^r \frac{\mathbf{H}(\rho)}{\rho^{2-\gamma_0}} d\rho = \frac{\mathbf{H}(\rho)}{\rho^{1-\gamma_0}} \Big|_0^r - \int_0^r \frac{d}{d\rho} \left(\frac{\mathbf{H}(\rho)}{\rho} \right) \rho^{\gamma_0} d\rho = \frac{\mathbf{H}(r)}{r^{1-\gamma_0}} - \int_0^r \frac{2\mathbf{E}(\rho)}{\rho^{1-\gamma_0}} d\rho. \quad (5.14)$$

The Cauchy–Schwarz inequality yields then the following bound for every ε :

$$|\mathbf{E}(r)| \leq \frac{\varepsilon}{r} \int_{\partial B_r} |\mathcal{X}|^2 + \frac{r}{4\varepsilon} \int_{\partial B_r} |D\mathcal{X}|^2 = \varepsilon \frac{\mathbf{H}(r)}{r} + \frac{r\mathbf{D}'(r)}{4\varepsilon}. \quad (5.15)$$

Therefore, by choosing $\varepsilon = \gamma_0/2$, we deduce (5.13) from (5.14) and (5.15). \square

Proof of Theorem 5.1. In view of Lemma 5.4, for r sufficiently small, the almost minimizing condition (4.4) reads as

$$\mathbf{D}(r) \leq C r \mathbf{D}'(r) + C \frac{\mathbf{H}(r)}{r^{1-\gamma_0}} + C \mathbf{m}_0^{1/2} r^{\gamma_0} \int_{\partial B_r} |\boldsymbol{\eta} \circ \mathcal{X}|.$$

Dividing by the radius and integrating we get

$$\begin{aligned} \int_0^r \frac{\mathbf{D}(s)}{s} ds &\leq C \int_0^r \left(\mathbf{D}'(\rho) + \frac{\mathbf{H}(\rho)}{\rho^{2-\gamma_0}} + \rho^{\gamma_0-1} \int_{\partial B_\rho} |\boldsymbol{\eta} \circ \mathcal{X}| \right) d\rho \\ &\stackrel{(5.13)}{\leq} C \mathbf{D}(r) + C \mathbf{F}(r) + C \mathbf{m}_0^{1/2} \int_{B_r} \frac{|\boldsymbol{\eta} \circ \mathcal{X}|}{|z|^{1-\gamma_0}} \\ &\stackrel{(2.14)}{\leq} C \mathbf{D}(r) + C (\boldsymbol{\Lambda}^{\gamma_0}(r) \mathbf{D}(r) + \mathbf{F}(r)) \leq C \mathbf{D}(r) + C r^{\gamma_0-1} \mathbf{H}(r). \end{aligned} \quad (5.16)$$

Therefore, using Lemma 5.2 we deduce that

$$\begin{aligned} \frac{\mathbf{H}(r)}{r} &= \int_0^r \frac{2\mathbf{E}(\rho)}{\rho} dt \stackrel{(5.4)}{\leq} C \int_0^r \frac{\mathbf{D}(\rho)}{\rho} d\rho \\ &\quad + C \int_0^r \left(\frac{\mathbf{H}(\rho)}{\rho^{2-2\gamma_0}} + \rho^{\gamma_0} \mathbf{D}'(\rho) + \rho^{\gamma_0} \frac{d}{d\rho} \|T - \mathbf{T}_F\|(\mathbf{p}^{-1}(\boldsymbol{\Psi}(B_\rho))) \right) d\rho \\ &\stackrel{(5.16)}{\leq} C \mathbf{D}(r) + C \frac{\mathbf{H}(r)}{r^{1-\gamma_0}} + C r^{\gamma_0} \mathbf{D}(r) + C \mathbf{F}(r) + C r^{\gamma_0} \|T - \mathbf{T}_F\|(\mathbf{p}^{-1}(\boldsymbol{\Psi}(B_r))) \\ &\stackrel{(2.15)\&(5.13)}{\leq} C \mathbf{D}(r) + C \frac{\mathbf{H}(r)}{r^{1-\gamma_0}}. \end{aligned}$$

For r sufficiently small this concludes the proof. \square

6. INNER VARIATIONS AND KEY ESTIMATES

Using the Poincaré inequality in Theorem 5.1, we can give very simple estimates of the error terms in the “inner variations” of the current T . The latter corresponds to deformations of T along appropriate vector fields which are tangent to \mathcal{M} . In order to state our main conclusion we need to introduce yet another quantity

$$\mathbf{G}(r) := \int_{\partial B_r} |D_\nu \mathcal{X}|^2. \quad (6.1)$$

Proposition 6.1 (Inner Variations). *There exist constants $C_{6.1} > 0$ and $\eta > 0$ such that, if $r > 0$ is small enough, then the following holds*

$$|\mathbf{D}'(r) - 2\mathbf{G}(r)| \leq C \mathcal{E}_{IV}(r), \quad (6.2)$$

where

$$\begin{aligned} \mathcal{E}_{IV}(r) &= r^{2\eta-1} \mathbf{D}(r) + \mathbf{D}(r)^\eta \mathbf{D}'(r) + \frac{\mathbf{m}_0^{1/2}}{r^{1-\gamma_0}} \int_{\partial B_r} |\boldsymbol{\eta} \circ \mathcal{X}(z, w)| \\ &\quad + \frac{d}{dr} \|T - \mathbf{T}_F\|(\mathbf{p}^{-1}(\Psi(B_r))). \end{aligned} \quad (6.3)$$

For further use we summarize in the next lemma a set of inequalities which will be used in the next sections and which are direct consequences of all the conclusions derived so far

Lemma 6.2. *There exist constant $C_{6.2} > 0$ and $\eta > 0$ such that for every r sufficiently small the following holds:*

$$\mathbf{F}(r) + r\mathbf{F}'(r) \leq C_{6.2} r^{\gamma_0} \mathbf{D}(r) \quad (6.4)$$

$$|\mathbf{L}(r)| \leq C_{6.2} r \mathbf{D}(r) \quad (6.5)$$

$$|\mathbf{L}'(r)| \leq C_{6.2} (\mathbf{H}(r) \mathbf{D}'(r))^{1/2} \quad (6.6)$$

$$\mathcal{E}_{OV} \leq C_{6.2} \mathbf{D}^{1+\eta}(r) + C_{6.2} \mathbf{F}(r) + C_{6.2} r \mathbf{D}^\eta(r) \mathbf{D}'(r) + C_{6.2} r \mathcal{E}_{BP}(r), \quad (6.7)$$

$$\mathcal{E}_{IV}(r) \leq C_{6.2} r^{2\eta-1} \mathbf{D}(r) + C_{6.2} \mathbf{D}(r)^\eta \mathbf{D}'(r) + C_{6.2} \mathcal{E}_{BP}(r), \quad (6.8)$$

where

$$\mathcal{E}_{BP}(r) := \frac{\mathbf{m}_0^{1/2}}{r^{1-\gamma_0}} \int_{\partial B_r} |\boldsymbol{\eta} \circ \mathcal{X}| + \frac{d}{dr} \|T - \mathbf{T}_F\|(\mathbf{p}^{-1}(\Psi(B_r)))$$

Moreover, for every $a > 0$ there exist constants $b_0(a), C(a) > 0$ such that

$$\mathbf{D}(r) \leq \frac{r \mathbf{D}'(r)}{2(2a+b)} + \frac{a(a+b) \mathbf{H}(r)}{r(2a+b)} + C(a) r \mathcal{E}_{IV}(r) \quad \forall b < b_0(a). \quad (6.9)$$

An important corollary of the previous lemma is the following

Corollary 6.3. *There exists a constant $C_{6.3} > 0$ such that, if η is the constant of Lemma 6.2, then for every $0 \leq \gamma < \eta$ and r sufficiently small, the nonnegative functions $\frac{\mathcal{E}_{IV}(r)}{r^\gamma \mathbf{D}(r)}$*

$\frac{\mathcal{E}_{OV}(r)}{r^{1+\gamma}\mathbf{D}(r)}$ are both integrable. Moreover, if we define the functions

$$\Sigma_{IV}(r) := \int_0^r \frac{\mathcal{E}_{IV}(s)}{s^\gamma \mathbf{D}(s)} ds, \quad (6.10)$$

$$\Sigma_{OV}(r) := \int_0^r \frac{\mathcal{E}_{OV}(s)}{s^\gamma \mathbf{D}(s)} ds, \quad (6.11)$$

$$\Sigma(r) := \Sigma_{IV}(r) + \Sigma_{OV}(r), \quad (6.12)$$

then

$$\Sigma(r) \leq C_{6.3} r^{\eta-\gamma}. \quad (6.13)$$

6.1. Proof of Proposition 6.1. We evaluate the first variation of T along a suitably defined vector field X . To this aim we fix a function $\phi \in C_c^\infty(]-1, 1[)$, symmetric, nonnegative and identically one on $]-1+1/j, 1-1/j[$ and with the property that $|\phi'| \leq Cj$. Then we introduce the vector field $Y: \mathcal{M} \rightarrow \mathbb{R}^{n+2}$ defined, for every $(z, w) \in \mathfrak{B} \setminus \{0\}$, by

$$Y(\Psi(z, w)) := \frac{|z|}{r} \phi\left(\frac{|z|}{r}\right) D_\nu \Psi(z, w) \in T_{\Psi(z, w)} \mathcal{M},$$

and extended to be 0 at the origin.

Next we define the vector field $X_i: \mathbf{V}_{a,u} \rightarrow \mathbb{R}^{n+2}$ by $X_i(p) := Y(\mathbf{p}(p))$. Note that X_i is the infinitesimal generator of a one parameter family of diffeomorphisms Φ_ε defined as $\Phi_\varepsilon(p) := \Gamma_\varepsilon(\mathbf{p}(p)) + p - \mathbf{p}(p)$, where Γ_ε is the one-parameter family of biLipschitz homeomorphisms of \mathcal{M} generated by Y . In fact, since Γ_ε fixes the origin, we can consider it as a C^{2,γ_0} map of $\mathcal{M} \setminus \{0\}$ onto itself. Note moreover that X_i is Lipschitz on the entire \mathfrak{B} .

Observe that, by Lemma 5.4 and the Poincaré inequality, $\mathbf{F}(r) \leq Cr^{\gamma_0} \mathbf{D}(r)$, so that $\mathbf{\Lambda}(r) \leq C \mathbf{D}(r)$. Moreover,

$$|D_{\mathcal{M}} Y|(\Psi(z, w)) + |\operatorname{div}_{\mathcal{M}} Y|(\Psi(z, w)) \leq -Cr^{-2} |z| \phi'\left(\frac{|z|}{r}\right) + Cr^{-1} \phi\left(\frac{|z|}{r}\right), \quad (6.14)$$

where we recall that $\phi' \leq 0$ on $[0, 1]$.

If r is small enough, by (2.13) we can argue as in [8, Section 3.3] and deduce that

$$\frac{1}{2} \left| \int_{\mathcal{M}} \left(|DN|^2 \operatorname{div}_{\mathcal{M}} Y - 2 \sum_{i=1}^Q \langle DN_i, (DN_i \cdot D_{\mathcal{M}} Y) \rangle \right) \right| \leq \sum_{k=1}^5 \operatorname{Err}_k^i,$$

where the error terms can be bounded in the following manner.

First of all,

$$\begin{aligned} \operatorname{Err}_1^i &= Q \left| \int_{\mathcal{M}} (\langle H_{\mathcal{M}}, \boldsymbol{\eta} \circ N \rangle \operatorname{div}_{\mathcal{M}} Y + \langle D_Y H_{\mathcal{M}}, \boldsymbol{\eta} \circ N \rangle) \right| \\ &\leq Cr^{-1} \mathbf{m}_0^{1/2} \int_{\mathfrak{B}} \left(\phi\left(\frac{|z|}{r}\right) |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{X}(z, w)| - \phi'\left(\frac{|z|}{r}\right) |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{X}(z, w)| \right) \\ &\stackrel{(2.14)}{\leq} Cr^{-1} \mathbf{D}^{1+\eta}(r) - C \mathbf{m}_0^{1/2} r^{\gamma_0-1} \int_{B_r} r^{-1} \phi'\left(\frac{|z|}{r}\right) |\boldsymbol{\eta} \circ \mathcal{X}(z, w)|, \end{aligned}$$

where in the first inequality we used (6.14) and the fact that

$$\langle D_Y H_{\mathcal{M}}, \boldsymbol{\eta} \circ N \rangle \leq |Y| |DH_{\mathcal{M}}| |\boldsymbol{\eta} \circ N| \leq C \frac{|z|}{r} \phi\left(\frac{|z|}{r}\right) |z|^{\gamma_0-2} |\boldsymbol{\eta} \circ \mathcal{X}|.$$

As for Err_2^i and Err_i^3 we have

$$\begin{aligned} \text{Err}_2^i &= C \int_{\mathcal{M}} |A_{\mathcal{M}}|^2 (|DY| |N|^2 + |Y| |N| |DN|) \\ &\leq C \mathbf{m}_0 \int_{\mathfrak{B}} \left[r^{-1} \left(-\frac{|z|}{r} \phi'\left(\frac{|z|}{r}\right) + \phi\left(\frac{|z|}{r}\right) \right) \frac{|\mathcal{X}|^2}{|z|^{2-2\gamma_0}} + \frac{|z|}{r} \phi\left(\frac{|z|}{r^2}\right) \frac{|\mathcal{X}| |D\mathcal{X}|}{|z|^{2-2\gamma_0}} \right] \\ &\leq C \mathbf{m}_0 r^{\gamma_0-1} \mathbf{D}(r) - C r^{-1} \int_{B_r} r^{-1} \phi'\left(\frac{|z|}{r}\right) \frac{|\mathcal{X}|^2}{|z|^{1-\gamma_0}}, \end{aligned}$$

and

$$\begin{aligned} \text{Err}_3^i &\leq C \int_{\mathcal{M}} \left(|Y| |A_{\mathcal{M}}| |DN|^2 (|N| + |DN|) + |DY| (|A_{\mathcal{M}}| |DN| |N|^2 + |DN|^4) \right) \\ &\leq C r^{\gamma_0-1} \mathbf{D}(r) - C \mathbf{D}(r)^\eta \int_{\mathfrak{B}} r^{-1} \phi'\left(\frac{|z|}{r}\right) |D\mathcal{X}|^2 + C r^{-1} \mathbf{D}(r)^\eta \int_{\mathfrak{B}} r^{-1} \phi\left(\frac{|z|}{r}\right) \frac{|\mathcal{X}|^2}{|z|^{2-\gamma_0}}. \end{aligned}$$

The errors Err_4^i and Err_5^i are the same as Err_4^o and Err_5^o respectively, in Section 5.2, evaluated along a different vector field. Proceeding in the same way as in the estimate of Err_4^o , we deduce

$$\begin{aligned} \text{Err}_4^i &= \int_{\text{spt}(T) \setminus \text{Im}(F)} |\text{div}_{\vec{T}} X_i| d\|T\| + \int_{\text{Im}(F) \setminus \text{spt}(T)} |\text{div}_{\vec{\mathbf{T}}_F} X_i| d\|\mathbf{T}_F\| \\ &\leq C r^{\gamma_0-1} \mathbf{D}(r) + C \underbrace{\int \alpha d\|T - \mathbf{T}_F\|}_{S(\phi)}. \end{aligned}$$

where $\alpha(p) = \varphi(\mathbf{p}(p))$ and $\varphi(\Psi(z, w)) = r^{-2} |z| \phi(r^{-1} |z|) - r^{-1} \phi'(r^{-1} |z|)$. In particular using (2.15) and the fact that $-\phi' \leq Cj$ on $[0, 1]$, we infer

$$S(\phi) \leq C r^{\gamma_0-1} \mathbf{D}(r) + C \frac{j}{r} \|T - \mathbf{T}_F\| (\mathbf{p}^{-1}(\Psi(B_r \setminus B_{r(1-1/j)}))).$$

As for Err_i^5 , we observe that it only appears in the cases (a) and (c) and arguing as in Section 5.2 we can bound it as

$$\text{Err}_i^5 \leq I_1 + \underbrace{\left| \int \langle X_i^\perp, h(\vec{\mathbf{T}}_F(p)) \rangle d\|\mathbf{T}_F\| \right|}_{I_2},$$

where $h(v_1 \wedge v_2) := \sum_{i=1}^2 A_\Sigma(v_i, v_i)$ and I_1 enjoys the same bounds as Err_i^4 . Following the argument of [8, Section 4.3] we can see that I_2 enjoys the same bounds as Err_i^1 and Err_i^2 .

To conclude the proof notice that, with analogous computation as in [4, Proposition 3.1],

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \int_{\mathcal{M}} |D(N \circ \Gamma_\varepsilon)|^2 = \int_{\mathcal{M}} \left(2 \sum_{i=1}^Q \langle DN_i : (DN_i \cdot D_{\mathcal{M}}Y) \rangle - |DN|^2 \operatorname{div}_{\mathcal{M}}Y \right). \quad (6.15)$$

However, by the conformal invariance of the Dirichlet energy, we have

$$\int_{\mathcal{M}} |D(N \circ \Gamma_\varepsilon)|^2 = \int_{\mathfrak{B}} |D(\mathcal{N} \circ \hat{\Gamma}_\varepsilon)|^2,$$

where $\hat{\Gamma}_\varepsilon$ is the one parameter family of diffeomorphisms generated by the vector field $\hat{Y} : \mathfrak{B} \rightarrow \mathfrak{B}$ defined by

$$\hat{Y}(z, w) := \frac{|z|}{r} \phi\left(\frac{|z|}{r}\right) \nu.$$

Hence

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \int_{\mathcal{M}} |D(N \circ \Gamma_\varepsilon)|^2 = \int_{\mathfrak{B}} \left(2 \sum_{i=1}^Q \langle D\mathcal{N}_i : (D\mathcal{N}_i \cdot D\hat{Y}) \rangle - |D\mathcal{N}|^2 \operatorname{div} \hat{Y} \right), \quad (6.16)$$

where the differentiation is taken with respect to the (local) flat structure of \mathfrak{B} .

In particular we conclude

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \int_{\mathcal{M}} |D(N \circ \Gamma_\varepsilon)|^2 = \int_{B_r} \frac{|z|}{r^2} \phi'\left(\frac{|z|}{r}\right) (2|D_\nu \mathcal{N}|^2 - |D\mathcal{N}|^2). \quad (6.17)$$

Collecting together (6.15), (6.17) and the error estimates, and letting ϕ converge to the to the indicator function of $[-1, 1]$ (namely letting $j \uparrow \infty$) we conclude the proof.

6.2. Proof of Lemma 6.2. The lemma is a very simple corollary of the estimates proven so far. (6.4) is a simple consequence of the Poincaré inequality (5.1) and of (5.13). Similarly, by Lemma 5.4, we have that $\mathbf{\Lambda}(r) \leq C \mathbf{D}(r)$, and therefore (6.7) follows in view of (6.4). The same arguments hold for (6.8). Next for (6.5) we can estimate as follows:

$$\begin{aligned} |\mathbf{L}(r)| &\leq C \mathbf{m}_0^{1/2} \int_{B_r} |\mathcal{N}| |D\mathcal{N}| \leq C \mathbf{m}_0^{1/2} \left(\int_0^r \mathbf{H}(t) dt \right)^{\frac{1}{2}} \mathbf{D}^{\frac{1}{2}}(r) \\ &\stackrel{(5.1)}{\leq} C \mathbf{m}_0^{1/2} \left(C_{5.1} \int_0^r t \mathbf{D}(t) dt \right)^{\frac{1}{2}} \mathbf{D}^{\frac{1}{2}}(r) \leq C \mathbf{m}_0^{1/2} r \mathbf{D}(r). \end{aligned} \quad (6.18)$$

Similarly

$$|\mathbf{L}'(r)| \leq C \mathbf{m}_0^{1/2} \int_{\partial B_r} |\mathcal{N}| |D\mathcal{N}| \leq C \mathbf{m}_0^{1/2} (\mathbf{D}'(r) \mathbf{H}(r))^{\frac{1}{2}}. \quad (6.19)$$

Finally, we notice that Proposition 6.1 implies

$$\left| \frac{\mathbf{D}'(r)}{2} - \int_{\partial B_r} |D_\tau \mathcal{N}|^2 \right| \leq C \mathcal{E}_{IV}(r).$$

Therefore, using the almost minimizing property in (4.5) and the Poincaré inequality we infer that

$$\mathbf{D}(r) \leq (1 + Cr) \left[\frac{r \mathbf{D}'(r)}{2(2a+b)} + \frac{a(a+b) \mathbf{H}(r)}{r(2a+b)} \right] + C(a)r \mathcal{E}_{IV}(r) + \mathcal{E}_{QM}(r) + Cr^{1+\sigma} \mathbf{D}'(r).$$

Absorbing the error term $r^{1+\sigma} \mathbf{D}'(r)$ and dividing by $(1 + Cr^\sigma)$ we get

$$\mathbf{D}(r) \leq \frac{r \mathbf{D}'(r)}{2(2a+b)} + \frac{a(a+b) \mathbf{H}(r)}{r(2a+b)} + C(a)r \mathcal{E}_{IV}(r) + \mathcal{E}_{QM}(r) + Cr^\sigma \mathbf{D}(r),$$

from which (6.9) follows straightforwardly by noticing that $\mathcal{E}_{QM}(r) + r \mathbf{D}(r) \leq Cr \mathcal{E}_{IV}(r)$.

6.3. Proof of Corollary 6.3. Recall first that $\eta < \gamma_0$. We start with $\mathcal{E}_{BP}(r)$. Notice that, using $\mathbf{H}(t) \leq Ct \mathbf{D}(t)$ together with the definition of $\mathbf{F}(r)$, we have

$$\int_0^r \left(\frac{1}{t^\gamma \mathbf{D}(t)} \right)' \mathbf{F}(t) dt \leq C \frac{\mathbf{F}(r)}{r^\gamma \mathbf{D}(r)} + C \int_0^r \frac{1}{t^\gamma \mathbf{D}(t)} \frac{\mathbf{H}(t)}{t^{2-\gamma_0}} dt \leq Cr^{\gamma_0-\gamma}$$

Next, by a simple integration by parts and the fact that $\mathbf{D}(r) \leq Cr^2$, we deduce

$$\begin{aligned} \int_0^r \frac{1}{t^\gamma \mathbf{D}(t)} \frac{d}{dt} \|T - \mathbf{T}_F\|(\mathbf{p}^{-1}(\Psi(B_t))) dt &= \frac{1}{r^\gamma \mathbf{D}(r)} \|T - \mathbf{T}_F\|(\mathbf{p}^{-1}(\Psi(B_r))) \\ &\quad + \int_0^r \left(\frac{1}{t^\gamma \mathbf{D}(t)} \right)' \|T - \mathbf{T}_F\|(\mathbf{p}^{-1}(\Psi(B_t))) dt \\ &\stackrel{(2.15)}{\leq} C \frac{\mathbf{D}^{1+\eta}(r) + \mathbf{F}(r)}{r^\gamma \mathbf{D}(r)} + \int_0^r \left(\frac{1}{t^\gamma \mathbf{D}(t)} \right)' (\mathbf{D}(t)^{1+\eta} + \mathbf{F}(t)) dt \leq Cr^{\eta-\gamma}. \end{aligned} \quad (6.20)$$

In a similar fashion we have

$$\begin{aligned} \int_0^r \frac{\mathbf{m}_0^{1/2}}{t^\gamma \mathbf{D}(t)} \int_{\partial B_t} \frac{|\boldsymbol{\eta} \circ \mathcal{X}(z, w)|}{t^{1-\gamma_0}} dt &\leq \frac{\mathbf{m}_0^{1/2}}{r^\gamma \mathbf{D}(r)} \int_{B_r} \frac{|\boldsymbol{\eta} \circ \mathcal{X}(z, w)|}{|z|^{1-\gamma_0}} \\ &\quad + \int_0^r \left(\frac{1}{t^\gamma \mathbf{D}(t)} \right)' \mathbf{m}_0^{1/2} \int_{B_t} \frac{|\boldsymbol{\eta} \circ \mathcal{X}(z, w)|}{|z|^{1-\gamma_0}} \\ &\stackrel{(2.14)}{\leq} C \frac{\mathbf{D}^{1+\eta}(r) + \mathbf{F}(r)}{r^\gamma \mathbf{D}(r)} + \int_0^r \left(\frac{1}{t^\gamma \mathbf{D}(t)} \right)' (\mathbf{D}(t)^{1+\eta} + \mathbf{F}(t)) dt \leq Cr^{\eta-\gamma}. \end{aligned} \quad (6.21)$$

so that

$$\int_0^r \frac{\mathcal{E}_{BP}(t)}{t^\gamma \mathbf{D}(t)} dt \leq Cr^{\eta-\gamma}$$

To conclude, we estimate separately the two nonnegative functions $\boldsymbol{\Sigma}_{IV}$ and $\boldsymbol{\Sigma}_{OV}$. In particular

$$\begin{aligned} \boldsymbol{\Sigma}_{IV}(r) &= \int_0^r \frac{\mathcal{E}_{IV}(t)}{t^\gamma \mathbf{D}(t)} dt \stackrel{(6.8)}{\leq} 2C_{6.2} \int_0^r \left(t^{\gamma_0-\gamma-1} + t^{-\gamma} \mathbf{D}(t)^{\eta-1} \mathbf{D}'(t) + \frac{\mathcal{E}_{BP}(t)}{t^\gamma \mathbf{D}(t)} \right) dt \\ &\leq Cr^{\eta-\gamma} (1 + \mathbf{D}(t)^{\eta/2}) \leq Cr^{\eta-\gamma}, \end{aligned} \quad (6.22)$$

where in the second inequality we used $\mathbf{D}(t) \leq C t^2$. Finally

$$\begin{aligned} \Sigma_{OV}(r) &= \int_0^r \frac{\mathcal{E}_{OV}(t)}{t^{1+\gamma} \mathbf{D}(t)} dt \\ &\stackrel{(6.7)}{\leq} C_{6.2} \int_0^r \left(\frac{\mathbf{D}^\eta(t)}{t^{1+\gamma}} + \frac{\mathbf{F}(t)}{t^{1+\gamma} \mathbf{D}(t)} + t^{-\gamma} \mathbf{D}^{\eta-1}(t) \mathbf{D}'(t) + \frac{\mathcal{E}_{BP}(t)}{t^\gamma \mathbf{D}(t)} \right) dt \\ &\leq C r^{\eta-\gamma}. \end{aligned} \tag{6.23}$$

7. ALMOST MONOTONICITY AND DECAY OF THE FREQUENCY FUNCTION

In this section we study the asymptotic behaviour of the normal approximation \mathcal{N} . The first step consists in proving approximate monotonicity and decay estimates for the frequency function.

For every $r \in (0, 1)$ such that $\mathbf{H}(r) > 0$, we set $\bar{\mathbf{I}}(r) := \frac{r \Omega(r)}{\mathbf{H}(r)}$ where we recall that

$$\Omega(r) := \begin{cases} \mathbf{D}(r) & \text{in the cases (a) and (b) of Definition 0.1;} \\ \mathbf{D}(r) + \mathbf{L}(r) & \text{in case (c).} \end{cases}$$

Furthermore we define $\bar{\mathbf{K}}(r) := \bar{\mathbf{I}}(r)^{-1}$ whenever $\Omega(r) \neq 0$. By (6.5) there exists $r_0 > 0$ such that

$$\frac{1}{2} \mathbf{D}(r) \leq (1 - Cr) \mathbf{D}(r) \leq \Omega(r) \leq (1 + Cr) \mathbf{D}(r) \leq 2 \mathbf{D}(r) \quad \forall r \leq r_0. \tag{7.1}$$

Having fixed r_0 , $\bar{\mathbf{K}}(r)$ is well defined whenever $\mathbf{D}(r) > 0$ and hence, by the Poincaré inequality, whenever $\bar{\mathbf{I}}(r)$ is defined. Moreover, if for some $\rho \leq r_0$ $\bar{\mathbf{K}}(\rho)$ is not well defined, that is $\Omega(\rho) = 0$, then obviously $\Omega(r) = \mathbf{D}(r) = 0$ for every $r \leq \rho$.

We are now ready to state the first important monotonicity estimate. From now on we assume of having fixed a γ

Theorem 7.1. *There exists a constant $C_{7.1} > 0$ with the following property: if $\mathbf{D}(r) > 0$ for some $r \leq r_0$, then (setting $\gamma = 0$ in (6.10) and (6.11)) the function*

$$\bar{\mathbf{K}}(r) \exp(-4r - 4\Sigma_{IV}(r)) - 4\Sigma_{OV}(r) \tag{7.2}$$

is monotone non-increasing on any interval $[a, b]$ where \mathbf{D} is nowhere 0. In particular, either there is $\bar{r} > 0$ such that $\mathbf{D}(\bar{r}) = 0$ or $\bar{\mathbf{K}}$ is well-defined on $]0, r_0[$ and the limit $K_0 := \lim_{r \rightarrow 0} \bar{\mathbf{K}}(r)$ exists.

A fundamental consequence of Theorem 7.1 is the following dichotomy.

Corollary 7.2. *There exists $\bar{r} > 0$ such that*

(A) *either $\bar{\mathbf{K}}(r)$ is well-defined for every $r \in]0, r_0[$, the limit*

$$K_0 := \lim_{r \downarrow 0} \bar{\mathbf{K}}(r) \tag{7.3}$$

is positive and thus there is a constant C and a radius \bar{r} such that

$$C^{-1} r \mathbf{D}(r) \leq \mathbf{H}(r) \leq C r \mathbf{D}(r) \quad \forall r \in]0, \bar{r}[; \tag{7.4}$$

(B) *or $T \llcorner \mathbf{p}^{-1}(\Psi(B_{\bar{r}})) = Q \llbracket \Psi(B_{\bar{r}}) \rrbracket$ for some positive \bar{r} .*

In turn, using the above dichotomy we will show

Theorem 7.3. *Assume that condition (i) in Theorem 2.8 fails. Then the frequency $\bar{\mathbf{I}}(r)$ is well-defined for every sufficiently small r and its limit $I_0 = \lim_{r \rightarrow 0} \bar{\mathbf{I}}(r) = K_0^{-1}$ exists and it is finite and positive. Moreover there exist constants $\lambda, C_{7.3}, H_0, D_0 > 0$ such that, for every r sufficiently small the following holds:*

$$|\mathbf{I}(r) - I_0| + \left| \frac{\mathbf{H}(r)}{r^{2I_0+1}} - H_0 \right| + \left| \frac{\mathbf{D}(r)}{r^{2I_0}} - D_0 \right| \leq C_{7.3} r^\lambda. \quad (7.5)$$

7.1. Proof of Theorem 7.1. In the first step we claim the monotonicity of the function $\bar{\mathbf{K}}(r) \exp(-\Sigma_{IV}(r)) - 2 \Sigma_{OV}(r)$ on any interval contained in $[a, b]$ on which \mathbf{D} is everywhere positive. Recalling that Ω and \mathbf{H} are absolutely continuous functions, we can compute the following derivative: for every $r \in [a, b]$

$$\begin{aligned} \bar{\mathbf{K}}'(r) &= \left(\frac{\mathbf{H}(r)}{r} \right)' \frac{1}{\Omega(r)} - \frac{\mathbf{H}(r)}{r} \frac{\Omega'(r)}{\Omega^2(r)} \\ &\stackrel{(5.3)}{\leq} \frac{1}{r\Omega^2(r)} (2\mathbf{E}(r)\Omega(r) - \mathbf{D}'(r)\mathbf{H}(r) + |\mathbf{L}'(r)|\mathbf{H}(r)). \end{aligned} \quad (7.6)$$

Then, either $\bar{\mathbf{K}}' \leq 0$, or the RHS of the inequality above is positive, that is

$$\mathbf{D}'(r)\mathbf{H}(r) \leq 2\mathbf{E}(r)\Omega(r) + |\mathbf{L}'(r)|\mathbf{H}(r) \stackrel{(6.6)}{\leq} 2\mathbf{E}(r)\Omega(r) + r\mathbf{D}'(r)\mathbf{H}(r) + \frac{\mathbf{H}^2(r)}{r}.$$

In turn, using $\mathbf{H}(r) \leq Cr\mathbf{D}(r) \leq Cr\Omega(r)$, the latter inequality implies

$$\mathbf{D}'(r)\mathbf{H}(r) \leq C\mathbf{E}(r)\Omega(r) + Cr\Omega^2(r). \quad (7.7)$$

From this we deduce

$$\mathbf{E}^2(r) \leq \mathbf{H}(r)\mathbf{G}(r) \leq \mathbf{H}(r)\mathbf{D}'(r) \leq C\Omega^2(r) + \frac{\mathbf{E}^2(r)}{2}$$

which implies that $\mathbf{E}(r) \leq C\Omega(r)$ and so, by (6.6),

$$|\mathbf{L}'(r)| \leq C\mathbf{m}_0^{1/2} (\mathbf{D}'(r)\mathbf{H}(r))^{1/2} \leq C\mathbf{m}_0^{1/2} \Omega(r). \quad (7.8)$$

Next using again the Cauchy-Schwarz inequality and (5.4), we have

$$\begin{aligned} \Omega(r)\mathbf{E}(r) &\leq \Omega(r)\mathbf{H}(r)^{1/2}\mathbf{G}(r)^{1/2} \leq \frac{\Omega(r)^2}{2} + \frac{\mathbf{H}(r)\mathbf{G}(r)}{2} \\ &\leq \frac{\Omega(r)\mathbf{E}(r)}{2} + \frac{\Omega(r)\mathcal{E}_{OV}(r)}{2} + \frac{\mathbf{H}(r)\mathbf{G}(r)}{2}, \end{aligned}$$

which implies

$$\Omega(r)\mathbf{E}(r) \leq \mathbf{H}(r)\mathbf{G}(r) + \Omega(r)\mathcal{E}_{OV}(r). \quad (7.9)$$

Collecting all these estimates together and using (6.2), we conclude that, if $\bar{\mathbf{K}}'(r) \geq 0$, then

$$\begin{aligned} \bar{\mathbf{K}}'(r) &\stackrel{(7.6)\&(7.9)}{\leq} \frac{1}{r\Omega^2(r)} (2\mathbf{H}(r)\mathbf{G}(r) - \mathbf{D}'(r)\mathbf{H}(r) + |\mathbf{L}'(r)|\mathbf{H}(r) + 2\Omega(r)\mathcal{E}_{OV}(r)) \\ &\stackrel{(6.2)\&(7.8)}{\leq} \frac{1}{r\Omega^2(r)} (2\mathbf{H}(r)\mathbf{G}(r) - 2\mathbf{H}(r)\mathbf{G}(r) + \Omega(r)\mathbf{H}(r) + \mathbf{H}(r)\mathcal{E}_{IV}(r) + 2\Omega(r)\mathcal{E}_{OV}(r)) \\ &\leq 2\frac{\mathcal{E}_{OV}(r)}{r\Omega(r)} + \bar{\mathbf{K}}(r) \left(1 + \frac{\mathcal{E}_{IV}(r)}{\Omega(r)}\right) \leq 4\frac{\mathcal{E}_{OV}(r)}{r\mathbf{D}(r)} + 4\bar{\mathbf{K}}(r) \left(1 + \frac{\mathcal{E}_{IV}(r)}{\mathbf{D}(r)}\right). \end{aligned} \quad (7.10)$$

On the other hand the final inequality

$$\bar{\mathbf{K}}'(r) \leq 4\frac{\mathcal{E}_{OV}(r)}{r\mathbf{D}(r)} + 4\bar{\mathbf{K}}(r) \left(1 + \frac{\mathcal{E}_{IV}(r)}{\mathbf{D}(r)}\right)$$

is certainly correct when $\bar{\mathbf{K}}'(r) \leq 0$, because the right hand side is positive. The monotonicity of the function in (7.2) is then obvious.

Next, as already observed, either \mathbf{D} is always positive, or it vanishes on some interval $]0, \bar{r}[$. If \mathbf{D} is always positive, then $\bar{\mathbf{K}}$ is well defined on $]0, r_0[$ and the existence of the limit $K_0 := \lim_{r \downarrow 0} \bar{\mathbf{K}}(r)$ is a direct consequence of (7.2) and Corollary 6.3.

7.2. Proof of Corollary 7.2. First of all observe that, if $\mathbf{D}(\bar{r})$ vanishes, then $\mathcal{N} \equiv Q \llbracket 0 \rrbracket$ on $B_{\bar{r}}$. In particular by (2.15) we conclude that we are in the alternative (B). We can thus assume, without loss of generality, that \mathbf{D} is positive on $]0, r_0[$. Assuming that K_0 vanishes we will then reach a contradiction.

Under the assumption $K_0 = 0$, consider the monotonicity of $\bar{\mathbf{K}}(r) \exp(-4\Sigma_{IV}(r)) - 4\Sigma_{OV}(r)$ between two radii $0 < s < r$ and let $s \rightarrow 0$ to get

$$\bar{\mathbf{K}}(r) \leq 4e^{4r+4\Sigma_{IV}(r)} \Sigma_{OV}(r) \leq C \Sigma_{OV}(r),$$

where the last inequality holds for r sufficiently small, since $\Sigma_{IV}(r) \leq Cr^\eta$ (recall that we have set $\gamma = 0$). Next observe that, since the function $\Sigma_{OV}(r)$ is non-decreasing,

$$\frac{\mathbf{F}(r)}{\mathbf{D}(r)} \leq \frac{1}{\mathbf{D}(r)} \int_0^r \frac{\mathbf{H}(s)\mathbf{D}(s)}{s^{2-\gamma_0}\mathbf{D}(s)} ds \leq C \int_0^r \frac{\bar{\mathbf{K}}(s)}{s^{1-\gamma_0}} ds \leq Cr^{\gamma_0} \Sigma_{OV}(r). \quad (7.11)$$

Moreover, integrating by parts:

$$\begin{aligned} &\int_0^r \frac{1}{\mathbf{D}(s)} \frac{d}{ds} \|T - \mathbf{T}_F\|(\mathbf{p}^{-1}(\Psi(B_s))) ds \\ &\stackrel{(2.15)}{\leq} C \frac{\mathbf{D}^{1+\eta}(r) + \mathbf{F}(r)}{\mathbf{D}(r)} + C \int_0^r \left(\frac{1}{\mathbf{D}(s)}\right)' (\mathbf{D}^{1+\eta}(s) + \mathbf{F}(s)) ds \\ &\leq C\mathbf{D}^\eta(r) + Cr^{\gamma_0}\Sigma_{OV}(r) + C\frac{\mathbf{F}(r)}{\mathbf{D}(r)} + C \int_0^r \frac{\mathbf{F}'(s)}{\mathbf{D}(s)} ds \\ &\leq C\mathbf{D}^\eta(r) + Cr^{\gamma_0}\Sigma_{OV}(r) + C \int_0^r \frac{\bar{\mathbf{K}}(s)}{s^{1-\gamma_0}} ds \leq C\mathbf{D}^\eta(r) + Cr^{\gamma_0}\Sigma_{OV}(r), \end{aligned} \quad (7.12)$$

where we have used repeatedly (7.1).

Using the latter in the formula for \mathcal{E}_{OV} we also conclude

$$\begin{aligned} & \Sigma_{OV}(r) \\ & \leq C \int_0^r \frac{1}{s\mathbf{D}(s)} \left(\mathbf{D}(s)^{1+\eta} + s\mathbf{D}^\eta(s)\mathbf{D}'(s) + \mathbf{F}(s) + s\frac{d}{ds}\|T - \mathbf{T}_F\|(\mathbf{p}^{-1}(\Psi(B_s))) \right) ds \\ & \leq C r^\eta \mathbf{D}(r)^{\eta/2} + C r^{\gamma_0} \Sigma_{OV}(r). \end{aligned}$$

Hence, for r sufficiently small,

$$\bar{\mathbf{K}}(r) \leq C \Sigma_{OV}(r) \leq C \mathbf{D}(r)^{\eta/2}. \quad (7.13)$$

In particular this implies that

$$\mathbf{H}(r) \leq C r \mathbf{D}(r)^{1+\eta/2}. \quad (7.14)$$

Combining this with (5.4) and the Cauchy-Schwarz inequality, we deduce

$$\begin{aligned} \frac{1}{2} \mathbf{D}(r) & \leq \Omega(r) \leq \frac{\mathbf{E}(r)}{r} + \mathcal{E}_{OV}(r) \leq \left(\frac{\mathbf{H}(r)}{r \mathbf{D}(r)^{\eta/4}} \right)^{1/2} (r \mathbf{D}'(r) \mathbf{D}(r)^{\eta/4})^{1/2} + \mathcal{E}_{OV}(r) \\ & \stackrel{(7.14)}{\leq} C \mathbf{D}(r)^{1+\eta/4} + C r \mathbf{D}(r)^{\eta/4} \mathbf{D}'(r) + \mathcal{E}_{OV}(r). \end{aligned}$$

Dividing the expression above by $r\mathbf{D}(r)$, integrating between two radii $0 < s < r$ and using the bound $\mathbf{D}(r) \leq C r^2$ we obtain

$$\log \left(\frac{r}{s} \right) \leq C \int_s^r \left(\frac{\mathbf{D}(\rho)^{\eta/4}}{\rho} + \mathbf{D}(\rho)^{\eta/4-1} \mathbf{D}'(\rho) + \frac{\mathcal{E}_{OV}(\rho)}{\rho \mathbf{D}(\rho)} \right) d\rho \leq C r^{\eta/2}.$$

Sending $s \rightarrow 0$ we get a contradiction.

7.3. Proof of Theorem 7.3. Clearly, if (i) in Theorem 2.8 does not hold, then \mathbf{D} is always positive and we are in alternative (A) of Corollary 7.2. Thus K_0 is positive and the first statement is obvious.

Let $\mathbf{K}(r) := \mathbf{I}(r)^{-1}$ and observe that by (7.1) we have

$$(1 - Cr)\mathbf{I}(r) \leq \bar{\mathbf{I}}(r) \leq (1 + Cr)\mathbf{I}(r), \quad \forall 0 \leq r \leq r_0,$$

which implies

$$(1 - Cr)\bar{\mathbf{K}}(r) \leq \mathbf{K}(r) \leq (1 + Cr)\bar{\mathbf{K}}(r) \quad \forall 0 \leq r \leq r_0,$$

so that in particular $\mathbf{K}(r) \leq C \bar{\mathbf{K}}(r) < \infty$ for every $0 < r < r_0$ and $\mathbf{K}(r) \rightarrow K_0$ as $r \rightarrow 0$. Using the monotonicity formula of Theorem 7.1 together with Corollary 6.3 we have

$$\bar{\mathbf{K}}(r) - K_0 \leq K_0(\exp(4r + 4\Sigma_{OV}(r)) - 1) + 4\Sigma_{IV}(r) \exp(4r + 4\Sigma_{OV}(r)) \leq C r^\eta. \quad (7.15)$$

Therefore

$$\mathbf{K}(r) - K_0 \leq C r^\eta + C \mathbf{K}(r) r \leq C r^\eta. \quad (7.16)$$

To control $\mathbf{K}(r) - K_0$ from below we apply (6.9) with $a = I_0 = \frac{1}{K_0}$ and $b = \lambda \leq \min\{\eta/2, b_0(I_0)\}$ to infer, after dividing by $r\mathbf{D}(r)$, that

$$-\frac{\mathbf{D}'(r)}{\mathbf{D}(r)} \leq \frac{2}{r} (I_0(I_0 + \lambda)\mathbf{K}(r) - (2I_0 + \lambda)).$$

Multiplying this expression by $\mathbf{K}(r) > 0$ and adding $2/r$, we get

$$\begin{aligned} \frac{2}{r} - \frac{\mathbf{D}'(r)}{\mathbf{D}(r)} \mathbf{K}(r) &\leq \frac{2}{r} \left[1 + I_0(I_0 + \lambda) \mathbf{K}^2(r) - (2I_0 + \lambda) \mathbf{K}(r) \right] + \frac{C \mathcal{E}_{IV}(r)}{\mathbf{D}(r)} \\ &\leq \frac{2}{r} I_0 \left(\mathbf{K}(r) - \frac{1}{I_0} \right) \left((I_0 + \lambda) \mathbf{K}(r) - 1 \right) + \frac{C \mathcal{E}_{IV}(r)}{\mathbf{D}(r)} \end{aligned} \quad (7.17)$$

Since $(I_0 + \lambda) \mathbf{K}(r)$ converges to $1 + \lambda K_0$, we easily deduce that for r small enough $(I_0 + \lambda) \mathbf{K}(r) - 1 \geq \frac{\lambda}{2} K_0$. Using this together with (7.16), we deduce from (7.17) that

$$\frac{2}{r} - \frac{\mathbf{D}'(r)}{\mathbf{D}(r)} \leq \frac{\lambda}{r} \left(\mathbf{K}(r) - \frac{1}{I_0} \right) + \frac{C \mathcal{E}_{IV}(r)}{\mathbf{D}(r)} + C \frac{r^\eta}{r}. \quad (7.18)$$

We next compute

$$\begin{aligned} \mathbf{K}'(r) &= \left(\frac{\mathbf{H}(r)}{r} \right)' \frac{1}{\mathbf{D}(r)} - \frac{\mathbf{H}(r)}{r \mathbf{D}(r)} \frac{\mathbf{D}'(r)}{\mathbf{D}(r)} \stackrel{(5.3)\&(7.1)}{\leq} \frac{2 \mathbf{E}(r)}{r \mathbf{D}(r)} - \frac{\mathbf{D}'(r)}{\mathbf{D}(r)} \mathbf{K}(r) \\ &\stackrel{(5.4)\&(7.1)}{\leq} \frac{2}{r} + C - \frac{\mathbf{D}'(r)}{\mathbf{D}(r)} \mathbf{K}(r) + C \frac{\mathcal{E}_{OV}(r)}{r \mathbf{D}(r)} \\ &\stackrel{(7.18)}{\leq} \frac{\lambda}{r} \left(\mathbf{K}(r) - \frac{1}{I_0} \right) + \frac{C \mathcal{E}_{IV}(r)}{\mathbf{D}(r)} + C \frac{\mathcal{E}_{OV}(r)}{r \mathbf{D}(r)} + C \frac{r^\eta}{r}. \end{aligned} \quad (7.19)$$

Recalling that $\mathbf{K}(r) \leq C$, we deduce

$$\frac{d}{dr} \left[\frac{\mathbf{K}(r) - K_0}{r^\lambda} \right] \leq C \frac{\mathcal{E}_{OV}(r)}{r^{1+\lambda} \mathbf{D}(r)} + C \frac{\mathcal{E}_{IV}(r)}{r^\lambda \mathbf{D}(r)} + C \frac{1}{r^{1+\lambda-\eta}}. \quad (7.20)$$

Integrating (7.20) on the interval $]s, r[$ and using (6.13), we get

$$\mathbf{K}(r) - K_0 \leq \frac{r^\lambda}{s^\lambda} (\mathbf{K}(s) - K_0) + C r^{\eta-\lambda}$$

that is $\mathbf{K}(s) - K_0 \geq -C s^\lambda$. The inequality $|\mathbf{K}(r) - K_0| \leq C r^\lambda$ easily implies $|\mathbf{I}(r) - I_0| \leq C r^\lambda$.

For what concerns the other inequalities we compute

$$\begin{aligned} \left[\log \left(\frac{\mathbf{H}(r)}{r^{2I_0+1}} \right) \right]' &= \frac{\mathbf{H}'(r)}{\mathbf{H}(r)} - \frac{2I_0 + 1}{r} = \frac{2 \mathbf{E}(r)}{r \mathbf{H}(r)} - \frac{2I_0}{r} \leq \frac{2 \mathbf{D}(r)}{\mathbf{H}(r)} - \frac{2I_0}{r} + C \frac{\mathcal{E}_{OV}(r)}{\mathbf{H}(r)} \\ &= \frac{2}{r} (\mathbf{I}(r) - I_0) + C \frac{\mathcal{E}_{OV}(r)}{\mathbf{H}(r)}, \end{aligned} \quad (7.21)$$

and similarly

$$\left[\log \left(\frac{\mathbf{H}(r)}{r^{2I_0+1}} \right) \right]' \geq \frac{2}{r} (\mathbf{I}(r) - I_0) - C \frac{\mathcal{E}_{OV}(r)}{\mathbf{H}(r)}. \quad (7.22)$$

Using that $|\mathbf{I}(r) - I_0| \leq C r^\lambda$, for r small enough we have the bound $r \mathbf{D}(r) \leq 2I_0 \mathbf{H}(r)$. Hence we can use (6.13) in the integrals of (7.21) and (7.22) to deduce the existence of the

limit

$$H_0 := \lim_{s \downarrow 0} \frac{\mathbf{H}(s)}{s^{2I_0+1}}, \quad \text{with} \quad \left| \frac{\mathbf{H}(r)}{r^{2I_0+1}} - H_0 \right| \leq C r^\lambda.$$

Moreover, from (7.21) we also infer that for r sufficiently small

$$H_0 \geq \frac{\mathbf{H}(r)}{r^{2I_0+1}} e^{-Cr^\lambda} > 0.$$

Finally the last assertion follows simply setting $D_0 := I_0 \cdot H_0$ and from

$$\begin{aligned} \left| \frac{\mathbf{D}(r)}{r^{2I_0}} - D_0 \right| &= \left| \mathbf{I}(r) \frac{\mathbf{H}(r)}{r^{2I_0+1}} - I_0 H_0 \right| \\ &\leq |\mathbf{I}(r) - I_0| \frac{\mathbf{H}(r)}{r^{2I_0+1}} + I_0 \left| \frac{\mathbf{H}(r)}{r^{2I_0+1}} - H_0 \right| \leq C r^\lambda. \end{aligned}$$

8. BLOW-UP AND PROOF OF THEOREM 2.8

As a consequence of the decay estimate in Theorem 7.3 we can show that suitable rescaling of the normal approximation N converge to a unique limiting profile. To this aim we consider for every $r \in (0, 1)$ the functions $f_r : \partial B_1 \rightarrow \mathcal{A}_{Q_1}(\mathbb{R}^{2+n})$ given by

$$f_r(z, w) := \frac{\mathcal{N}(i_r(z, w))}{r^{I_0}},$$

where we recall that $i_r(z, w) = (rz, r^{1/\bar{Q}}w)$. We recall also that $T_0\mathcal{M} = \mathbb{R}^2 \times \{0\}$, and $T_0\Sigma = \mathbb{R}^2 \times \mathbb{R}^{\bar{n}} \times \{0\}$. In the following, with a slight abuse of notation, we write $\mathbb{R}^{\bar{n}}$ for the subspace $\{0\} \times \mathbb{R}^{\bar{n}} \times \{0\}$.

The final step in the proof of Theorem 2.8 is then the following proposition.

Proposition 8.1. *Assume alternative (i) in Theorem 2.8 fails and let I_0 and λ be the positive numbers of Theorem 7.3. Then $I_0 > 1$ and there exists a function $f_0 : \partial B_1 \rightarrow \mathcal{A}_Q(\mathbb{R}^{\bar{n}})$ such that*

- (i) $\boldsymbol{\eta} \circ f_0 = 0$ and $f_0 \not\equiv Q_1 \llbracket 0 \rrbracket$;
- (ii) for every r sufficiently small

$$\mathcal{G}(f_r(z, w), f_0(z, w)) \leq C r^{\lambda/16} \quad \forall (z, w) \in \partial B_1; \quad (8.1)$$

- (iii) the I_0 -homogeneous extension $g(z, w) := |z|^{I_0} f_0\left(\frac{z}{|z|}, \frac{w}{|w|}\right)$ is nontrivial and Dir-minimizing.

In particular, by (iii) $\text{Im}(g) \setminus \{0\} \subset \mathbb{R}^{2+n}$ is a real analytic submanifold.

Theorem 2.8 follows immediately from Proposition 8.1 and Theorem 7.3.

Proof of Theorem 2.8. Since we have identified $\mathbb{R}^{\bar{n}}$ with $\{0\} \times \mathbb{R}^{\bar{n}} \times \{0\}$, it is obvious that the map g has all the properties claimed in (ii), namely it is Dir-minimizing, $\boldsymbol{\eta} \circ g \equiv 0$ and it is nontrivial. (2.16) is a corollary of (8.1) provided $a_0 \leq \frac{\lambda}{16}$. Next note that (2.18) has

been shown in Theorem 5.1. As for (2.17) observe that, if $4\rho \leq r < 1$, then, by Theorem 7.3,

$$D_0(r - 2\rho)^{2I_0} - C(r - 2\rho)^{2I_0 + \lambda} \leq \mathbf{D}(r - 2\rho) \leq \mathbf{D}(r + 2\rho) \leq D_0(r + 2\rho)^{2I_0} + C(r + 2\rho)^{2I_0 + \lambda}.$$

Since $2I_0 > 2$, (2.17) follows easily from

$$\int_{B_{r+2\rho} \setminus B_{r-2\rho}} |D\mathcal{X}|^2 = \mathbf{D}(r + 2\rho) - \mathbf{D}(r - 2\rho),$$

provided $a_0 \leq \lambda$. \square

The rest of this final section of the note is devoted to the proof Proposition 8.1, which is split in several steps. Before starting with it, let us however observe that the conclusion $I_0 > 1$ is an obvious consequence of the decay estimates of Theorem 7.3 and the fact that $\mathbf{D}(r) \leq Cr^{2+2\gamma_0}$.

8.1. Step 1: uniqueness of the limit f_0 . For r sufficiently small and $s \in [\frac{r}{2}, r]$, we start estimating the following quantity:

$$\int_{\partial B_1} \mathcal{G}(f_r, f_s)^2 \leq (r - s) \int_{\partial B_1} \int_s^r \left| \frac{d}{dt} f_t(z, w) \right|^2 dt. \quad (8.2)$$

Using the differentiability properties of Lipschitz multiple valued functions and the 1-dimensional theory in [4, Section 1.1.2] (note that $t \mapsto \mathcal{X}(i_t(z, w))$ is a Lipschitz map), we easily infer that

$$\begin{aligned} \left| \frac{d}{dt} f_t(z, w) \right|^2 &= \sum_{j=1}^Q \left| \frac{D\mathcal{X}_j(i_t(z, w)) \cdot z}{t^{I_0}} - I_0 \frac{\mathcal{X}_j(i_t(z, w))}{t^{I_0+1}} \right|^2 \\ &= \frac{|z|^2 |\partial_{\hat{r}} \mathcal{X}|^2(i_t(z, w))}{t^{2I_0}} - 2I_0 \frac{|z|}{t^{2I_0+1}} \sum_{j=1}^Q \langle \partial_{\hat{r}} \mathcal{X}_j, \mathcal{X}_j \rangle(i_t(z, w)) + \frac{|\mathcal{X}|^2(i_t(z, w))}{t^{2I_0+2}}. \end{aligned}$$

Therefore, by the change of variable $(z', w') = i_t(z, w)$ in (8.2) we infer that

$$\begin{aligned} \int_{\partial B_1} \mathcal{G}(f_r, f_s)^2 &\leq \frac{r}{2} \int_{r/2}^r \left(\frac{\mathbf{G}(t)}{t^{2I_0+1}} - 2I_0 \frac{\mathbf{E}(t)}{t^{2I_0+2}} + I_0^2 \frac{\mathbf{H}(t)}{t^{2I_0+3}} \right) dt \\ &\leq \frac{r}{2} \int_{r/2}^r \left(\frac{\mathbf{D}'(t)}{2t^{2I_0+1}} - 2I_0 \frac{\mathbf{D}(t)}{t^{2I_0+2}} + I_0^2 \frac{\mathbf{H}(t)}{t^{2I_0+3}} + C \frac{\mathcal{E}_{IV}(t)}{t^{2I_0+1}} + C \frac{\mathcal{E}_{OV}(t)}{t^{2I_0+2}} \right) dt \\ &= \frac{r}{2} \int_{r/2}^r \left[\frac{1}{2t} \left(\frac{\mathbf{D}(t)}{t^{2I_0}} \right)' + I_0 \frac{\mathbf{H}(t)}{t^{2I_0+3}} (I_0 - \mathbf{I}(t)) + C \frac{\mathcal{E}_{IV}(t)}{t^{2I_0+1}} + C \frac{\mathcal{E}_{OV}(t)}{t^{2I_0+2}} \right] dt. \end{aligned}$$

Using Theorem 7.3, we can then conclude that

$$\begin{aligned} \int_{\partial B_1} \mathcal{G}(f_r, f_s)^2 &\leq C \left| \frac{\mathbf{D}(r)}{r^{2I_0}} - \frac{\mathbf{D}(\frac{r}{2})}{(\frac{r}{2})^{2I_0}} \right| + C \int_{r/2}^r \left[\frac{|I_0 - \mathbf{I}(t)|}{t} + C \frac{\mathcal{E}_{IV}(t)}{\mathbf{D}(t)} + C \frac{\mathcal{E}_{OV}(t)}{t \mathbf{D}(t)} \right] dt \\ &\leq Cr^\lambda. \end{aligned} \quad (8.3)$$

By an elementary dyadic argument analogous to that of [4, Theorem 5.3], we then infer the existence of $f_0 : \partial B_1 \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+n})$ such that, for r sufficiently small,

$$\|\mathcal{G}(f_r, f_0)\|_{L^2(\partial B_1)}^2 \leq C r^\lambda. \quad (8.4)$$

8.2. Step 2: uniform convergence. Set next $h(z, w) := \mathcal{G}\left(\frac{\mathcal{N}(z, w)}{|z|^{I_0}}, \frac{\mathcal{N}(i_{1/2}(z, w))}{|z/2|^{I_0}}\right)$. It follows from (8.3) that for r sufficiently small

$$\int_{B_r} h^2 \leq \int_0^r \int_{\partial B_1} \mathcal{G}(f_t, f_{t/2})^2 t dt \stackrel{(8.3)}{\leq} C r^{2+\lambda}, \quad (8.5)$$

and from (2.12) and (2.13)

$$\text{Lip}(h|_{B_1 \setminus B_s}) \leq C s^{-I_0}. \quad (8.6)$$

Moreover, for every $\rho < |z|/4$ we claim the estimate

$$\int_{B_\rho(z, w)} |Dh|^2 \leq C \rho + C |z|^\lambda. \quad (8.7)$$

Indeed $|Dh| \leq C \left| D \left(\frac{\mathcal{N}}{|z|^{I_0}} \right) \right|$ and by Theorem 7.3

$$\begin{aligned} \int_{B_\rho(z, w)} \left| D \left(\frac{\mathcal{N}}{|z|^{I_0}} \right) \right|^2 &\leq 2 \int_{|z|-\rho}^{|z|+\rho} \int_{\partial B_t} \left(\frac{|D\mathcal{N}|^2}{t^{2I_0}} + I_0^2 \frac{|\mathcal{N}|^2}{t^{2I_0+2}} \right) dt \\ &\leq 2 \int_{|z|-\rho}^{|z|+\rho} \left(\left(\frac{\mathbf{D}(t)}{t^{2I_0}} \right)' + 2 I_0 \frac{\mathbf{D}(t)}{t^{2I_0+1}} + I_0^2 \frac{\mathbf{H}(t)}{t^{2I_0+2}} \right) dt \\ &\leq C (|z| + \rho)^\lambda + C \log \left(\frac{|z| + \rho}{|z| - \rho} \right) \leq C |z|^\lambda + C \frac{\rho}{|z|}. \end{aligned}$$

In particular, applying (8.5), (8.6) and (8.7) with $\rho = |z|^{1+\lambda/4}$, we infer that for every point $p = (z, w) \in \mathfrak{B}_{\bar{Q}}$ with $|z|$ sufficiently small

$$\begin{aligned} h(p) &\leq \left| h(p) - \fint_{B_{\frac{|z|^{1+\lambda/4}}{2^k}}(p)} h \right| + \sum_{i=0}^{k-1} \left| \fint_{B_{\frac{|z|^{1+\lambda/4}}{2^i}}(p)} h - \fint_{B_{\frac{|z|^{1+\lambda/4}}{2^{i+1}}}(p)} h \right| + \fint_{B_{|z|^{1+\lambda/4}}(p)} h \\ &\leq \text{Lip}(h|_{B_1(p) \setminus B_{|z|/2}(p)}) \frac{|z|^{1+\lambda/4}}{2^k} + C \sum_{i=0}^{k-1} \frac{|z|^{1+\lambda/4}}{2^i} \fint_{B_{\frac{|z|^{1+\lambda/4}}{2^i}}(p)} |Dh| + \fint_{B_{|z|^{1+\lambda/4}}(p)} h \\ &\stackrel{(8.6)}{\leq} C |z|^{1+\lambda/4} + C \sum_{i=0}^{k-1} \left(\int_{B_{\frac{|z|^{1+\lambda/4}}{2^i}}(p)} |Dh|^2 \right)^{\frac{1}{2}} + \frac{C}{|z|^{1+\lambda/4}} \left(\int_{B_{2|z|}} |h|^2 \right)^{\frac{1}{2}}, \quad (8.8) \end{aligned}$$

where we have used the standard Poincaré inequality

$$\left| \fint_{B_r} f - \fint_{B_{\frac{r}{2}}} f \right| \leq C r \fint_{B_r} |Df| \quad f \in W^{1,2}.$$

Now choose $k \in \mathbb{N}$ such that $\frac{|z|^{1+\lambda/4}}{2^k} < |z|^{1+\lambda/4+I_0} \leq \frac{|z|^{1+\lambda/4}}{2^{k-1}}$ (in particular $k \leq |\log |z||$) and use (8.5) together with (8.7) to bound

$$h(z, w) \leq C |z|^{1+\lambda/4} + C |\log |z|| |z|^{\lambda/8} + C |z|^{\lambda/4} \leq C |z|^{\lambda/16}, \quad (8.9)$$

This gives that, for a sufficiently small r ,

$$\max_{\partial B_1} \mathcal{G}(f_r, f_{r/2}) \leq C r^{\lambda/16}.$$

Thus

$$\max_{\partial B_1} \mathcal{G}(f_r, f_0) \leq \sum_{k=0}^{\infty} \mathcal{G}(f_{r2^{-k}}, f_{r2^{-k-1}}) \leq C r^{\lambda/16}.$$

8.3. Step 3: nontriviality of the limit and other properties. To show that $f_0 \neq \mathcal{Q} \llbracket 0 \rrbracket$ it is enough to observe that, by Theorem 7.3,

$$\int_{\partial B_1} |f_0|^2 = \lim_{r \rightarrow 0} \int_{\partial B_1} |f_r|^2 = \lim_{r \rightarrow 0} \frac{\mathbf{H}(r)}{r^{2I_0+1}} = H_0 > 0.$$

In order to show that $\boldsymbol{\eta} \circ f_0 \equiv 0$, we notice that by a simple slicing argument combined with (2.14) there exists a sequence of radii $r_k \in [2^{-k-1}, 2^{-k}]$ such that

$$\begin{aligned} \int_{\partial B_{r_k}} |\boldsymbol{\eta} \circ \mathcal{X}| &\leq 2^{k+1} \int_{B_{2^{-k}} \setminus B_{2^{-k-1}}} |\boldsymbol{\eta} \circ \mathcal{X}| \leq C r_k^{\gamma_0} \int_{B_{2^{-k}}} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{X}| \\ &\leq C r_k^{\gamma_0+2\eta} \mathbf{D}(2r_k) \leq C r_k^{\gamma_0+2\eta+2I_0}, \end{aligned} \quad (8.10)$$

from which

$$\begin{aligned} \int_{\partial B_1} |\boldsymbol{\eta} \circ f_0| &= \lim_{r_k \rightarrow 0} \int_{\partial B_1} |\boldsymbol{\eta} \circ f_{r_k}| = \lim_{r_k \rightarrow 0} r_k^{-I_0-1} \int_{\partial B_{r_k}} |\boldsymbol{\eta} \circ \mathcal{X}| \\ &\leq C \lim_{r_k \rightarrow 0} r_k^{\gamma_0+2\eta+I_0-1} = 0. \end{aligned}$$

Next we show that f_0 takes values in $\mathbb{R}^{\bar{n}}$. We start by showing that f_0 must take values in $T_0\Sigma = \mathbb{R}^{2+\bar{n}} \times \{0\}$. Indeed, if we set $f_r(z, w) := \bar{\mathcal{X}}(i_r(z, w))$, using (A.3) and $|\mathcal{X}|(i_r(z, w)) \leq C r^{1+\gamma_0/2}$ we conclude

$$\int_{\partial B_1} \mathcal{G}(f_r, \bar{f}_r)^2 \leq \frac{C r^2}{r^{2I_0+1}} \int_{\partial B_r} |\mathcal{X}|^2 \leq C r^2,$$

which implies that $f_0(z, w) \in \mathcal{A}_{\mathcal{Q}}(T_0\Sigma)$.

Next observe that $f_r(z, w) = \sum_i \llbracket \mathcal{X}_i(i_r(z, w)) \rrbracket$ has the property that each $\mathcal{X}_i(i_r(z, w))$ is orthogonal to $T_{\Psi(i_r(z, w))} \mathcal{M}$. In particular, if $|z| = 1$ and $r \downarrow 0$, the tangent planes $T_{\Psi(i_r(z, w))} \mathcal{M}$ converge to $\mathbb{R}^2 \times \{0\}$: it follows, by the uniform convergence of f_r to f_0 , that $f_0(z, w) = \sum_i \llbracket (f_0)_i(z, w) \rrbracket$ for some $(f_0)_i(z, w)$ which are orthogonal to $\mathbb{R}^2 \times \{0\}$. We thus conclude that each $(f_0)_i(z, w)$ belongs to $\{0\} \times \mathbb{R}^{\bar{n}} \times \{0\}$.

8.4. Step 4: Minimality of g . In order to complete the proof of Proposition 8.1 we need to show that g is Dir-minimizing. Given the homogeneity of g in the radial direction, it suffices to show that there is no $W^{1,2}$ multifunction $h : B_1 \rightarrow \mathcal{A}_Q(\mathbb{R}^{\bar{n}})$ which has the same trace of g on ∂B_1 and less energy on B_1 . Assume thus by contradiction that there is an $h \in W^{1,2}(B_1, \mathcal{A}_Q(\mathbb{R}^{\bar{n}}))$ such that $h|_{\partial B_1}$ and

$$\int |Dh|^2 \leq \int |Dg|^2 - \delta \quad (8.11)$$

for some positive $\delta > 0$. Recall the definition of $W^{1,2}$ according to Remark 2.7: using the map \mathbf{W} in there and the functions $h \circ \mathbf{W}$ and $g \circ \mathbf{W}$ we can use the theory of [4] and assume that $h \circ \mathbf{W}$ is a Dir-minimizer on the euclidean disk $D_1 \subset \mathbb{R}^2$. Observe also that, since $\boldsymbol{\eta} \circ g \equiv 0$, we must have $\boldsymbol{\eta} \circ h \equiv 0$ as well. Indeed since $h \circ \mathbf{W} = g \circ \mathbf{W}$ on ∂D_1 , we have $\boldsymbol{\eta} \circ h \circ \mathbf{W} = \boldsymbol{\eta} \circ g \circ \mathbf{W} = 0$ on the boundary and considering that

$$\int_{D_1} \sum_i |D(h_i \circ \mathbf{W} - \boldsymbol{\eta} \circ h \circ \mathbf{W})|^2 \leq \int_{D_1} |D(h \circ \mathbf{W})|^2 - Q \int_{D_1} |D(\boldsymbol{\eta} \circ h \circ \mathbf{W})|^2,$$

the minimality of $h \circ \mathbf{W}$ forces the Dirichlet energy of $\boldsymbol{\eta} \circ h \circ \mathbf{W}$ to vanish identically.

Using (2.14), the decay $\mathbf{D}(r) \leq C r^{2I_0}$ and a Fubini-type argument we can find a sequence of radii $s_j \rightarrow 0$ such that

$$\int_{\partial B_1} |Df_0|^2 \leq \limsup_j \int_{\partial B_1} |Df_{s_j}|^2 \leq \limsup_j \frac{\mathbf{D}'(s_j)}{s_j^{2I_0-1}} \leq C. \quad (8.12)$$

We now wish to “smooth” h , i.e. to approximate it with a sequence of Lipschitz maps h_ε such that $\boldsymbol{\eta} \circ h_\varepsilon \equiv 0$,

$$\int_{B_1} |Dh_\varepsilon|^2 - |Dh|^2 \leq \varepsilon^2 \quad (8.13)$$

$$\int_{\partial B_1} \mathcal{G}(f_0, h_\varepsilon)^2 + \left| \int_{\partial B_1} |Df_0|^2 - |Dh_\varepsilon|^2 \right| \leq \varepsilon^2. \quad (8.14)$$

We would like to appeal to [6, Lemma 3.5], but there is the slight technical complication that \mathfrak{B} is not regular. We postpone this technical step and continue with the argument assuming the existence of the approximations h_ε .

Next we would like to apply [6, Lemma 3.6] to h_ε and $\mathbf{p}_{T_0\Sigma}(f_{s_j}) =: \bar{f}_{s_j}$, to get a family of competitor functions $(\hat{f}_{s_j}) \subset W^{1,2}(B_1, \mathcal{A}_Q(\mathbb{R}^{2+\bar{n}}))$, such that $\hat{f}_{s_j}|_{\partial B_1} = \bar{f}_{s_j}|_{\partial B_1}$ and

$$\int_{B_1} |D\hat{f}_{s_j}|^2 \leq \int_{B_1} |Dh_\varepsilon|^2 + \varepsilon \int_{\partial B_1} (|D_\tau h_\varepsilon|^2 + |D_\tau \bar{f}_{s_j}|^2) + \frac{C}{\varepsilon} \int_{\partial B_1} \mathcal{G}(h_\varepsilon, \bar{f}_{s_j})^2, \quad (8.15)$$

$$\text{Lip}(\hat{f}_{s_j}) \leq C \left(\text{Lip}(h_\varepsilon) + \text{Lip}(\bar{f}_{s_j}) + \frac{1}{\varepsilon} \sup_{\partial B_1} \mathcal{G}(\bar{f}_{s_j}, h_\varepsilon) \right) \quad (8.16)$$

$$\boldsymbol{\eta} \circ \hat{f}_{s_j} = \boldsymbol{\eta} \circ \bar{f}_{s_j}. \quad (8.17)$$

Again, this is not straightforward because [6, Lemma 3.6] is stated for euclidean domains. We postpone this second technical problem and continue with our argument assuming the existence of \hat{f}_{s_j} .

We are now ready to define our competitor function. We set $\bar{\mathcal{L}}_{s_j}(z, w) := s_j^{I_0} \hat{f}_{s_j}(i_{1/s_j}(z, w))$ and, observing that $\bar{\mathcal{L}}_{s_j}$ takes value in $\mathcal{A}_Q(T_0\Sigma)$, we use (3.1) to define a corresponding \mathcal{L}_{s_j} , which clearly is a competitor \mathcal{N} in B_{s_j} according to Definition 3.1. Moreover

$$\text{Lip}(\mathcal{L}_{s_j}) \leq C s_j^{I_0+1} \text{Lip}(\hat{f}_{s_j}|_{B_1}) \stackrel{(8.6)}{\leq} C s_j^\eta.$$

Therefore we can apply Proposition 3.2 with $\bar{\mathcal{L}} = \bar{\mathcal{L}}_{s_j}$. In particular, taking into account Theorem 7.3 and (8.12), we conclude that

$$\int_{B_{s_j}} |D\bar{\mathcal{N}}|^2 \leq (1 + C s_j) \int_{B_{s_j}} |D\bar{\mathcal{L}}_{s_j}|^2 + C m_0^{1/2} \int_{B_{s_j}} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{L}_{s_j}| + C s_j^{2I_0+\eta}.$$

Next, recall the inequality (4.6):

$$\int_{B_{s_j}} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \mathcal{L}_{s_j}| \leq C \int_{B_{s_j}} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \bar{\mathcal{L}}_{s_j}| + C \int_{B_{s_j}} |z|^{\gamma_0-1} |\bar{\mathcal{L}}_{s_j}|^2.$$

By (8.17) the first term in the right hand side equals indeed

$$C \int_{B_{s_j}} |z|^{\gamma_0-1} |\boldsymbol{\eta} \circ \bar{\mathcal{N}}| \leq C s_j^\eta \mathbf{D}(s_j) \leq C s_j^{2I_0+\eta}.$$

For the second term we use the Poincaré inequality

$$\int_{B_{s_j}} |z|^{\gamma_0-1} |\bar{\mathcal{L}}_{s_j}|^2 \leq C s_j^{1+\gamma_0} \int_{B_{s_j}} |D\bar{\mathcal{L}}_{s_j}|^2 + C s_j^{\gamma_0} \int_{\partial B_{s_j}} |\bar{\mathcal{L}}_{s_j}|^2, \quad (8.18)$$

whose proof will be given in Lemma A.1.

Using that

$$\int_{\partial B_{s_j}} |\bar{\mathcal{L}}_{s_j}|^2 = \int_{\partial B_{s_j}} |\bar{\mathcal{N}}|^2 = \mathbf{H}(s_j) \leq C s_j^{2I_0+1},$$

we easily conclude that

$$\int_{B_{s_j}} |D\bar{\mathcal{N}}|^2 \leq (1 + C s_j) \int_{B_{s_j}} |D\bar{\mathcal{L}}_{s_j}|^2 + C s_j^{2I_0+\eta}. \quad (8.19)$$

Changing variables and dividing by $s_j^{2I_0}$ we infer that

$$\int_{B_1} |D\bar{f}_{s_j}| \leq \int_{B_1} |D\hat{f}_{s_j}|^2 + C s_j^\eta. \quad (8.20)$$

Using (8.13), (8.14) and (8.15), we conclude

$$\begin{aligned} \int_{B_1} |D\bar{f}_{s_j}|^2 &\leq \int_{B_1} |Dh|^2 + Cs_j^\eta + C\varepsilon + \frac{C}{\varepsilon} \int_{\partial B_1} \mathcal{G}(f_0, \bar{f}_{s_j})^2 \\ &\leq \int_{B_1} |Dg|^2 - \delta + Cs_j^\eta + C\varepsilon + \frac{C}{\varepsilon} \int_{\partial B_1} \mathcal{G}(f_0, \bar{f}_{s_j})^2, \end{aligned}$$

where the constant C is independent of ε . In particular, if we fix ε sufficiently small and we then let $s_j \downarrow 0$, by the uniform convergence of f_{s_j} to f_0 on ∂B_1 we conclude

$$\limsup_{j \rightarrow \infty} \int_{B_1} |D\bar{f}_{s_j}|^2 \leq \int_{B_1} |Dg|^2 - \frac{\delta}{2}.$$

Since however $f_{s_j} \rightarrow g$ in B_1 , the latter inequality contradicts the semicontinuity of the Dirichlet energy.

8.5. Step 5: Technical leftovers. First of all we show the existence of the map h_ε as in (8.13) and (8.14). We consider $h \circ \mathbf{W}$, which is defined on the closed unit disk $\bar{D}_1 \subset \mathbb{R}^2$. We then can apply [6, Lemma 3.5] to the latter map and generate approximations \hat{h}_ε which satisfy the bounds (8.13) and (8.14) with D_1 in place of B_1 and $h \circ \mathbf{W}$ in place of h . The maps $h_\varepsilon := \hat{h}_\varepsilon \circ \mathbf{W}$ would then satisfy the desired estimates because of the conformality of \mathbf{W}^{-1} (which keeps the Dirichlet energy invariant) and its regularity in $B_1 \setminus \{0\}$ (which results into the loss of a constant factor in (8.14)). However the resulting map would not be Lipschitz because of the singularity of \mathbf{W}^{-1} in the origin. To overcome this difficulty it suffices to perturb slightly \hat{h}_ε so that it is constant in a small neighborhood of the origin. As for the condition $\boldsymbol{\eta} \circ h_\varepsilon \equiv 0$, this can easily be achieved subtracting the average to whichever extension satisfies (8.13) and (8.14).

Secondly we show the existence of \hat{f}_{s_j} . First of all we observe that the condition (8.17) can be easily achieved after we prove the existence of a map which satisfies the other two conditions: as above it suffices to subtract the average of this map and add back $\boldsymbol{\eta} \circ \bar{f}_{s_j}$. At this point we observe that it suffices, as above, to compose with the map \mathbf{W} , apply [4, Lemma 2.14] and [5, Lemma 3.6] and compose the resulting map with \mathbf{W}^{-1} : indeed the latter would coincide with $h_\varepsilon \circ \mathbf{W}$ on $D_{1-\varepsilon}$ and on the complement \mathbf{W}^{-1} is regular.

APPENDIX A. SOME USEFUL LEMMAS.

The first lemma is a simple version of the Poincaré inequality for $W^{1,2}$ functions.

Lemma A.1. *There exists a universal constant $C > 0$ such that the following two inequalities hold for every $f \in W^{1,2}(B_r, \mathcal{A}_Q)$ with $B_r \subset \mathfrak{B}_Q$:*

$$\int_{B_r} |f|^2 \leq Cr^2 \int_{B_r} |Df|^2 + Cr \int_{\partial B_r} |f|^2 \tag{A.1}$$

$$\int_{B_r} |z|^{\gamma_0-1} |f|^2 \leq Cr^{1+\gamma_0} \int_{B_r} |Df|^2 + Cr^{\gamma_0} \int_{\partial B_1} |f|^2. \tag{A.2}$$

Proof. By approximation we can assume, without loss of generality, that f is Lipschitz and, by scaling, it suffices to show the inequalities (A.1) and (A.2) on the ball B_1 . Fixing $|z| = 1$ and integrating along rays

$$|f(rz, r^{1/Q}w)|^2 \leq 2|f(z, w)|^2 + 2 \int_r^1 |Df(tz, t^{1/Q}w)|^2 dt.$$

Using radial coordinates we then conclude

$$\int_{B_1} |z|^{\gamma_0-1} |f|^2 \leq C \int_{\partial B_1} |f|^2 + \int_{\partial B_1} \int_0^1 r_0^\gamma \int_r^1 |Df(tz, t^{1/Q}w)|^2 dt dr dz.$$

Using Fubini the latter integral can be rewritten as

$$\int_0^1 \int_{\partial B_1} |Df(tz, t^{1/Q}w)|^2 \int_0^t r^{\gamma_0} dt dz dr \leq \int_0^1 t \int_{\partial B_1} |Df(tz, t^{1/Q}w)|^2 dz dr.$$

This completes the proof of (A.2). The proof of (A.1) is a simple variation of this one and is left to the reader. \square

Lemma A.2. *Let $\bar{\mathcal{L}}: \mathfrak{B}_{\bar{Q}} \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+\bar{n}})$ be Lipschitz and consider the map $\mathcal{L}: \mathfrak{B}_{\bar{Q}} \rightarrow \mathcal{A}_Q(\mathbb{R}^{2+n})$ defined by (3.1). Then there exists a constant $C := C(\|\Psi_0\|_{C^3}) > 0$ such that*

$$\mathcal{G}(\mathcal{L}, \bar{\mathcal{L}})(z, w) \leq Cr |\bar{\mathcal{L}}|(z, w) + C |\bar{\mathcal{L}}|^2(z, w), \quad \forall (z, w) \in B_r \quad (\text{A.3})$$

$$\int_{B_r} |D\mathcal{L}|^2 \leq (1 + Cr) \int_{B_r} |D\bar{\mathcal{L}}|^2 + Cr \int_{\partial B_r} |\bar{\mathcal{L}}|^2. \quad (\text{A.4})$$

Proof. For what concerns (A.3), observe that $D\Psi(0) = 0$ implies $\|D\Psi\|_{L^\infty(B_r)} \leq Cr$. Therefore, by the C^3 regularity of Ψ , we get

$$\begin{aligned} \mathcal{G}(\mathcal{L}, \bar{\mathcal{L}})(z, w) &= \sum_{j=1}^Q |\Psi(\mathbf{p}_0(\Psi) + \bar{\mathcal{L}}_j) - \Psi(\mathbf{p}_0(\Psi))|(z, w) \\ &\leq \|D\Psi\|(\Psi(z, w)) |\bar{\mathcal{L}}|(z, w) + \|A_\Sigma\| |\bar{\mathcal{L}}|^2(z, w) \\ &\leq Cr |\bar{\mathcal{L}}|(z, w) + C |\bar{\mathcal{L}}|^2. \end{aligned}$$

An analogous computation gives

$$\int_{B_r} |D\mathcal{L}|^2 \leq (1 + Cr) \int_{B_r} |D\bar{\mathcal{L}}|^2 + C \int_{B_r} |\bar{\mathcal{L}}|^2$$

and we conclude (A.4) using Lemma A.1. \square

REFERENCES

- [1] Frederick J. Almgren, Jr. *Almgren's big regularity paper*, volume 1 of *World Scientific Monograph Series in Mathematics*. World Scientific Publishing Co. Inc., River Edge, NJ, 2000.
- [2] Costante Bellettini and Tristan Rivière. The regularity of special Legendrian integral cycles. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 11(1):61–142, 2012.
- [3] Sheldon Xu-Dong Chang. Two-dimensional area minimizing integral currents are classical minimal surfaces. *J. Amer. Math. Soc.*, 1(4):699–778, 1988.

- [4] Camillo De Lellis and Emanuele Spadaro. Q -valued functions revisited. *Mem. Amer. Math. Soc.*, 211(991):vi+79, 2011.
- [5] Camillo De Lellis and Emanuele Spadaro. Multiple valued functions and integral currents. *Preprint*, 2013.
- [6] Camillo De Lellis and Emanuele Spadaro. Regularity of area-minimizing currents I: gradient L^p estimates. *Preprint*, 2013.
- [7] Camillo De Lellis and Emanuele Spadaro. Regularity of area-minimizing currents II: center manifold. *Preprint*, 2013.
- [8] Camillo De Lellis and Emanuele Spadaro. Regularity of area-minimizing currents III: blow-up. *Preprint*, 2013.
- [9] Camillo De Lellis, Emanuele Spadaro, and Luca Spolaor. Regularity theory for 2-dimensional almost minimal currents II: branched center manifold. 2015.
- [10] Camillo De Lellis, Emanuele Spadaro, and Luca Spolaor. Regularity theory for 2-dimensional almost minimizing currents I: Lipschitz approximation. 2015.
- [11] Camillo De Lellis, Emanuele Spadaro, and Luca Spolaor. Uniqueness of tangent cones for 2-dimensional almost minimizing currents. 2015.
- [12] David Pumberger and Tristan Rivière. Uniqueness of tangent cones for semicalibrated integral 2-cycles. *Duke Math. J.*, 152(3):441–480, 2010.
- [13] Tristan Rivière and Gang Tian. The singular set of J -holomorphic maps into projective algebraic varieties. *J. Reine Angew. Math.*, 570:47–87, 2004.
- [14] Leon Simon. *Lectures on geometric measure theory*, volume 3 of *Proceedings of the Centre for Mathematical Analysis, Australian National University*. Australian National University Centre for Mathematical Analysis, Canberra, 1983.
- [15] Brian White. Tangent cones to two-dimensional area-minimizing integral currents are unique. *Duke Math. J.*, 50(1):143–160, 1983.
- [16] Anton Zorich. Flat surfaces. In *Frontiers in number theory, physics, and geometry. I*, pages 437–583. Springer, Berlin, 2006.